

PUBLIC HEALTH REPORTS

VOL. 41

SEPTEMBER 10, 1926

No. 37

THE RADIOACTIVITY OF NATURAL WATERS

By W. D. COLLINS, Chemist in Charge, Quality of Water Division, United States Geological Survey

Every natural water is more or less radioactive, and therefore the recognition of the presence of a small amount of radioactive material in any spring water does not set that water apart from other natural waters.

Knowledge of the presence of radioactive substances in waters is comparatively recent. Therefore, the radioactivity has been seized as something to talk about and advertise as a remarkable and unique property of many waters which are no more unique in their radioactivity than they are in their wetness.

Physicians and others who have given thought to the subject have long recognized that, in general, much better results are obtained from the use of mineral waters at their sources than from the use of the same waters after they have been shipped in bottles. The common explanation has been that at resorts more water is taken, and at the same time the diet and other living conditions of the patient are better regulated than at home. Even without special medical attention at the resort there is usually rest, recreation, and freedom from the normal cares of life.

When radioactivity was first studied and it was found that many famous medicinal waters contained radium emanation, this fact was immediately taken as an explanation of the greater benefits derived from the use of the waters at their sources. The radium emanation is half gone in about 3.8 days after the water has been taken from its source and practically all gone within 30 days. So far the explanation seems reasonable. The first determinations of radioactivity of natural waters were nearly all made on samples from well-known springs. Later studies brought out great differences in the radioactivity of waters that seemed to produce identical beneficial effects and also showed radioactivity to be a universal property of natural water. As the use of radium emanation in the treatment of disease was developed, it appeared that the doses necessary to produce detectable effects could not be obtained by drinking any reasonable quantity of one of the naturally radioactive spring waters; of most waters it would be necessary to drink from 100 to 1,000 gallons a day.

In *New and Nonofficial Remedies*, 1925, page 308, it is stated that the Council on Pharmacy and Chemistry of the American

Medical Association will not accept any radium solution for internal use the dosage of which is less than 2 micrograms a day, or any radium emanation generator which yields less than 2 microcuries of emanation during each 24 hours.

To obtain the dose of 2 micrograms by drinking 1 gallon of water, which is considerably more than most people drink in a day, the radioactivity of the water would have to be about 500 millimicrocuries¹ per liter. As long ago as 1913 Lazarus, in his *Handbuch der Radium-Biologie und-Therapie*, stated (p. 200) that, in the use of waters having less than 20 millimicrocuries of radioactivity per liter, the radioactivity probably has no appreciable part in the beneficial effects obtained. It thus appears that the radioactivity should be between 20 and 500 millimicrocuries per liter in order to be considered at all as a factor in the use of a water as medicine, and that even within these limits the radioactivity is not likely to be the controlling factor.

In 1905 Boltwood² reported the radioactivity of waters from a group of 44 springs which have been used medicinally for bathing or drinking. The radioactivity ranged from 0.016 to 8.8 millimicrocuries per liter and only two had over 2.5 millimicrocuries per liter. Half of the results were between 0.1 and 0.5. Schlundt and Moore³ found less than 0.5 millimicrocurie per liter for 40 out of 88 springs in Yellowstone National Park; the maximum reported was 2.68 millimicrocuries per liter.

Measurements by the writer quoted by Skinner and Sale⁴ in a discussion of the radioactivity of miscellaneous waters examined in the Bureau of Chemistry gave results ranging from 0.007 to 1.1 millimicrocuries per liter for 15 samples of spring waters collected in 1911, 1912, and 1914 in Virginia, Wisconsin, and Massachusetts. Some of these samples were taken from widely known springs.

The text by Lazarus noted above contains an article by Sommer which gives results of measurement of radioactivity of 422 waters from springs in Germany. The radioactivity of 72 per cent of these waters was reported as less than 2.5 millimicrocuries per liter.

Moore and Whittemore⁵ measured the radioactivity of 14 waters at Saratoga Springs, N. Y., and found from 0.039 to 0.88 millimicrocurie per liter. Satterly and Elworthy⁶ report the radioactivity of waters from 23 wells and 47 springs in Canada. The results range from

¹ A millimicrocurie is the radioactivity corresponding to one-billionth of a gram of radium (0.000000001 gram). This unit is used in discussions of the radioactivity of natural waters because results expressed in it are numbers of moderate size. * Some reports have been made in units of one-tenth to one-thousandth of a millimicrocurie, but all the results quoted in the present discussion have been converted to millimicrocuries.

² *Am. Jour. Sci.*, 4th ser., vol. 20, pp. 128-32, 1905.

³ *U. S. Geol. Survey Bull.* 395, 1909.

⁴ *Jour. Ind. and Eng. Chem.*, vol. 14, pp. 949-950, 1922.

⁵ *Jour. Ind. and Eng. Chem.*, vol. 6, pp. 552-553, 1914.

⁶ Canada Dept. Mines, Mines Branch, Bull. 16, 1917.

0.0014 to 0.176 millimicrocurie per liter for the well waters and from 0.0112 to 0.64 for the spring waters.

Measurements by O. C. Lester of the radioactivity of 178 mineral waters of Colorado are given in Bulletin 11 of the State Geological Survey, published in 1920. The results range from "none" to 30.5 millimicrocuries per liter. The report "none" signifies only that no radioactivity was detected with the apparatus used and does not indicate the complete absence of radioactivity. Of the waters with measurable radioactivity 85 per cent had less than 2.5 millimicrocuries per liter, 14 per cent from 2.5 to 5, and 6 per cent from 5 to 30.

Other results quoted by the authors cited above serve to show the almost universal radioactivity of natural waters as they occur in the ground and the exceedingly small quantities of radioactivity in even the most radioactive of these waters when compared with the quantities which those who have studied the subject consider necessary to produce any therapeutic effect.

The best available evidence based on scientific studies of the treatment of disease with radium emanation, on measurements of radioactivity of natural spring waters, and on the reported uses of the spring waters, leads to the conclusion that, up to this time, it has not been shown that the small amounts of radioactivity found in natural waters have any effect on the medicinal value of the waters.

THE PHYSIOLOGICAL EFFECTS OF CURRENTS OF VERY HIGH FREQUENCY (135,000,000 TO 8,300,000 CYCLES PER SECOND)

By J. W. SCHERESCHEWSKY, Surgeon, United States Public Health Service

This paper reports the results of studies of the effects upon small laboratory animals (mice) of electrical oscillations of very high frequency generated by a vacuum tube oscillator.

The modern development of the vacuum tube oscillator and associated circuits permits the generation of continuous wave currents of relatively pure wave form of very high frequency, sharply emitted at the frequency to which the circuit is tuned. This is not the case in the usual type of high frequency apparatus used for therapeutic purposes. Here, the oscillations are produced by condenser discharge across a spark gap. The oscillations produced in this way have a large decrement; the wave form is impure, giving rise to many harmonics; the emitted wave is broad, and consistent operation at the frequencies worked with in the studies here reported is difficult to obtain. Moreover, the oscillations generated are in the form of discontinuous trains separated by a period during which the up-building of energy for the next train occurs.

The physiological effects of high frequency currents generated by the vacuum tube oscillator appear to have been but little investigated.

The only reference found by a search through the literature was a report by Gosset, Gutmann, Lakhovsky, and Magrou¹ on the effects of very high frequency radiation emitted by a vacuum tube oscillator upon plant tumors caused in the geranium by *Bacterium tumefaciens*.

The authors report that three geranium plants, bearing tumors caused by inoculation with the above-mentioned organism, were exposed to radiations emitted by a vacuum-tube oscillator at a frequency said to be about 150,000,000 cycles per second. One plant was given 2 exposures of three hours on consecutive days, one plant 3, and one plant 11 such exposures. After 16 days from the first exposure, the tumors, after growing in the interval, began suddenly to necrose. The necrotic process was said to be complete in about 15 days, so that the tumor could be detached by slight traction. In 16 control plants the tumors grew rapidly to enormous size and recurred after surgical excision. Details, however, as to the apparatus and method of exposure are lacking.

In the studies here reported the first step was a study of oscillators which might generate oscillations of sufficiently high frequency and the development of suitable auxiliary circuits in which the effects of such currents on small laboratory animals (mice) might be investigated.

At this point it is desired to express grateful acknowledgment to Prof. George W. Pierce and E. L. Chaffee, of the Cruft High Tension Laboratory of Harvard University, for permitting the use of the facilities of the Cruft High Tension Laboratory in the preliminary work and for much helpful advice; also to Mr. M. L. Dow and Mr. F. H. Drake, of the same laboratory, for collaboration, valuable assistance, and advice in the working-out, construction, and setting-up of apparatus and in making preliminary runs and tests.

Further grateful acknowledgment is made to Prof. M. J. Rosenau, of the Department of Preventive Medicine and Hygiene of the Harvard Medical School, for the use of the facilities of his department in the subsequent work. It is also desired to thank the General Electric Co. for furnishing three special vacuum tubes of low internal capacity.

Description of Apparatus

Vacuum tube oscillator.—After considerable preliminary work in testing the suitability of various types of oscillators, the following types of circuit were settled upon as satisfactory:

For the generation of the highest frequency currents employed, (i. e., from 200,000,000 cycles to 60,000,000 cycles per second) the type of oscillator described by Huxford² was found excellently

¹ A. Gosset, A. Gutmann, G. Lakhovsky, and I. Magrou: Essai de Therapeutique du "Cancer Experimental" des Plantes. Comptes Rendus de la Société de Biologie, Vol. 91, 1924, pp. 626-628.

² Huxford, W. S.: Standing Waves on Parallel Wires. Physical Review, Corning, N. Y., 2nd Series, Vol. 25, 1925, pp. 696-695.

adapted. In the design of circuits intended to oscillate at these high frequencies, the interelectrode capacity of the tube, negligible at low frequencies, becomes an important limiting factor on the frequency of the oscillations generated. In Huxford's circuit the capacity formed by the tube elements is in series with the variable tuning capacity. Because of well-known physical considerations, this has the effect of reducing the minimum capacity at which the oscillating circuit may be resonated, and thus permits the generation of frequencies




FIG. 1.—Circuit of oscillators Nos. 1 and 2 (Huxford circuit)

- L_1 , Grid inductance
- L_2 , Plate inductance
- C_1 , Tuning condenser
- RFC, Radio-frequency choke-coils
- MA, Plate milliammeter
- AM, Filament ammeter
- R, Variable resistance (10,000-200,000 ohms)

higher than would be possible with a tube having a given interelectrode capacity, were this capacity in shunt with the tuning capacity as is the case, for instance, in the well-known Hartley circuit.

Two oscillators of the type described by Huxford were constructed, one having a range of from about 200,000,000 cycles to 85,000,000 cycles and the other with a range of 150,000,000 cycles to 60,000,000 cycles per second. Figure 1 shows the circuit network and the accompanying photographs show the appearance of the oscillators.

Reference to the diagram and photograph of oscillator No. 1 (Pl. I) shows that the tuning inductance consists essentially of two parallel brass rods 24 centimeters long, 4.7 millimeters in diameter, and spaced 5 centimeters, supported at each end by vertical bakelite strips. At the vacuum tube end the upper rod is connected to the grid, while the lower is connected to the plate of the tube. This lower rod is revolvably mounted and serves as a support for the rotary plate of the tuning condenser which is mounted on the rod by means of a sleeve and set-screw, so that it may be moved and fixed at any point on the lower rod. The upper rod, which is not revolvable, carries the stator portion of the tuning condenser. This is suspended from a small brass block, bored to a sliding fit upon the upper rod and secured by a set-screw so that it may be appropriately located with respect to the rotary plate. The stator portion of the condenser consists of two plates spaced 3.1 millimeters from each other, with a radius of 4.7 centimeters. The rotary plate has a radius of 3.75 centimeters.

This arrangement of the tuning condenser which permits it to be slid to different sites along the rods is an important means of varying the oscillator frequency, as the range obtained by varying the condenser alone, at any one position on the rods is rather narrow. To get the benefit of the full range of frequencies of which the oscillator is capable, it is necessary to alter the value of the inductance in the oscillating circuit. This is done by sliding the rotor and stator portions of the condenser to different sites along the rods, the inductance in the circuit being a maximum when the condenser is slid as far away from the tube as possible. The farther ends of the rods are connected through suitable choke-coils to the filament and to the plate battery, respectively, the upper rod being connected to the negative filament through a variable resistance. The resistance used was a Bradley, variable between the values of 10,000 and 200,000 ohms. In the circuit, as described by Huxford, no resistance was employed; but in this instance its use appeared to add stability and efficiency, besides, because of its biasing action, reducing consumption of plate current and undue heating of the plate.

As shown in the diagram, the oscillating current was confined to the oscillating circuit by the use of suitable choke coils. These were four in number and were located, one in each leg of the filament, one at the outer end of the lower rod between it and the plate supply, and one at the outer end of the upper rod between this and the grid-biasing resistance. In oscillator No. 1 these choke coils consisted of 23 turns of No. 20 D. C. C. wire wound in a spiral 1.25 centimeters in diameter, each turn being slightly spaced from the next. They were readily made by winding the required number of turns tightly around a $\frac{1}{2}$ -inch rod and then slipping them off, the small diameter of the coil and the natural stiffness of the wire rendering

the coils self-supporting. While 23 turn coils were used and found to work well, this number is by no means critical. It could probably be varied several turns in either direction without changing results.

Oscillator No. 2 was a duplicate, in constructional details, of the first oscillator, except for larger dimensions. The rods were 38 centimeters long and spaced 11.5 centimeters instead of 5 centimeters. The tuning condenser, too, was larger, having five instead of three plates. Because of the wider spacing of the rods, it was necessary to make the brass block which carried the stator of the condenser considerably longer than in the smaller model.

It will be noted, from the photograph, that, in the smaller oscillator, no socket for the tube was used, the tube being mounted by inserting the base, until stopped by the pin, in a hole of proper diameter, in a horizontally mounted bakelite strip. The plate, grid, and filament leads were then soldered directly to their respective tube prongs. This was done to avoid introducing unnecessary capacity in the circuit through the use of a socket. In the other oscillators, however, where lower frequencies were generated, this precaution was needless; consequently, for the sake of convenience, a socket was employed.

The vacuum tube used for the generation of high-frequency current was the UX 210. This tube has a thoriated filament, is rated at 7.5 watts, is used with a maximum filament current of 1.25 amperes at 7.5 volts. The plate potential employed was 400 volts, furnished by a suitable number of 6,000 m. a.-hour lead storage-cells.

This tube was found to oscillate vigorously in the circuit just described up to a frequency of 158,000,000 cycles. At this point the internal capacity of the tube and the inductance of its leads were so great that oscillations of higher frequency could not be obtained. However, by the use of the special tubes (shown in Plate II) having a low internal capacity, which were obtained from the research laboratories of the General Electric Co., with the smaller oscillator it was possible to generate effective oscillations of a frequency around 200,000,000 cycles per second, and feebler ones of possibly 230,000,000 cycles.

As shown by the photograph, this tube has the filament and plate leads brought out at the lower end of the tube while the grid lead is brought out at the top. This reduces the capacity between leads within the tube which, in the ordinary type of mounting, may be several micromicrofarads.

Measurements of the internal capacity of this tube made at 1,000 cycles on the Croft Laboratory capacity bridge showed this to be between 3 and 4 micromicrofarads, less than one-half that of a standard UV 202, 5-watt tube.

While the tubes, when received, were provided with bases, these were removed, by heating, before use so as to reduce the tube capacity to a minimum.

It was found that the generation of these high frequencies was very hard on the tubes, it being necessary, in order to secure adequate output, to increase the filament and plate currents considerably beyond the normal standard. This resulted in burn-outs, cracking of seals, and, in the case of the UX 210 tube, of depletion of electron emission beyond the possibility of reactivation.






FIG. 2.—Circuit diagram of oscillator No. 3 (Hartley circuit,

- L_1 , Plate and grid inductance
- C_1 , Tuning condenser (5-plate and 17-plate)
- C_2 , Blocking condenser
- R_1 , Variable resistance (10,000-200,000 ohms)
- R. F. C., Radio-frequency choke-coil
- M_1 , Plate milliammeter
- M_2 , Filament ammeter


For frequencies of 60,000,000 cycles per second and less it was no longer found necessary to use the type of oscillator just described, which, though reliable and stable in operation, nevertheless, because of limited range was not as well suited to the exploration of the lower frequencies, as the ever useful and efficient Hartley circuit. Consequently, for these lower frequencies, the latter circuit with parallel plate feed, through a choke coil, was employed as shown in Figure 2 and the accompanying photograph.




Oscillator No. 1 (Huxford circuit). Range 230,000,000 to 85,000,000 cycles




Oscillator No. 2 (Huxford circuit). Range 150,000,000 to 60,000,000 cycles




Oscillator No. 3 (Hartley circuit). Range 50,000,000 to 23,000,000 cycles




Oscillator No. 3 (Hartley circuit). Range 20,000,000 to 7,250,000 cycles



Special 5-watt vacuum tube with low internal capacity



Auxiliary-coupled circuit showing mouse holder, thermocouple, and microammeter



For the inductance, single turn loops of No. 14 to No. 4 wire were used according to size and the need for rigidity. These loops, as shown in the photograph, were supported on the tuning condenser (a low-priced 5-plate affair with bakelite end plates which was found to function as satisfactorily as one of the most expensive "low loss" construction).

For frequencies below 12,000,000 cycles, however, a high-grade 17-plate condenser was used. The plate-blocking condenser, a Dubilier micadon, had the value of .00025 mfd. The filament return was taken off the mid-point of the tuning inductance, the connection to the negative filament being made, as in the case of the other oscillators, through a variable Bradley resistance, the usual resistance values used being from 10,000 to 15,000 ohms.

Choke coils of several sizes were inserted in the plate feed, larger choke coils being used for the lower frequencies. As in the case of the oscillators previously described, these consisted of a self-supporting spiral of No. 18 or No. 20 D. C. C. wire, the number of turns varying from 23 to 65 or 70 and the diameter from 2.5 to 4 centimeters, the turns being spaced about the diameter of the wire.

With this type of oscillator, simply by varying the diameter of the single turn used as the inductance, with the 5-plate condenser it was possible to go from 66,000,000 cycles per second (using a 6-centimeter loop) to 10,000,000 cycles per second (using a turn of No. 4 copper wire 76 centimeters in diameter). With a single turn loop 32 centimeters in diameter tuned by a .00035 mfd. condenser, frequencies as low as 7,500,000 cycles could be readily generated.

In all the oscillators the filament temperature was controlled by a 2-ohm rheostat. A Weston ammeter in the filament circuit and a Weston 0-300 milliammeter in the plate circuit indicated the filament and plate currents, respectively. The latter instrument was placed in the negative plate-battery lead and was protected by the insertion of a short strip of $\frac{1}{4}$ -ampere fuse wire.

It was generally found that, for adequate output, the necessary plate current was considerably greater at the higher than at the lower frequencies. At the highest frequencies, during some of the runs, plate current readings of from 100-120 milliamperes were not unusual, but at the lower frequencies from 50 to 80 milliamperes was the usual value.

Frequency determination.—For the purpose of ascertaining the frequency at which the oscillator was operating, use was made of a Lecher parallel wire system. This consisted of two No. 12 parallel copper wires 7.5 centimeters apart and 11 meters long, stretched from standards 29 centimeters above the level of the workbench. Turn-buckles at one end of the wire system served to tighten the wires.

The slider was a rectangular piece of 22-gauge brass plate 12.5 by 8 centimeters, provided with two parallel slots 7 centimeters long and 7.5 centimeters apart to enable it to engage the wires. Supporting sliders of bakelite, grooved to fit the wire and attached front and rear steadied the slider plate, enabling it to travel smoothly. A waxed string passing over pulleys attached to the standards allowed the slider to be moved to any position from the operator's location. A plumb bob suspended by a waxed thread from the slider indicated its position with respect to a wave-length scale laid out on the work-bench beneath. Resonance of the wire system with the oscillator frequency was indicated by a Weston thermogalvanometer connected across the oscillator end of the wires.

In a system of this character, as is well known, standing waves are formed on the parallel wire system in a series one-half wave in length between nodes. If the slider is placed at one of these nodes the wave will travel along the wires to the slider and be reflected back, the total distance traveled being equal to the distance between wave crests, and therefore corresponding to the wave length. When the slider is located in this fashion at a node the meter will show the maximum deflection.

The system is readily calibrated by setting the oscillator at some frequency sufficiently high to permit the development of several successive nodes on the wire system, moving the slider to each of these points in succession and measuring the interval between them; and averaging the measurements which will be found to differ from each other by less than a centimeter. In this way orientation points on the wave-length scale are readily located from which the wave-length scale may be laid out, the wave-length measured, and hence the frequency determined.

Although, owing to the capacity between the wires and also to the surface of the bench, a slight error is introduced, so that the apparent frequency is slightly higher than the actual, still this method of measuring the wave length, and consequently the frequency, is remarkably accurate, the error probably being one-half of 1 per cent or less.

Since available space permitted a wire system only 11 meters long, wave lengths only up to approximately 21 meters, or frequencies somewhat less than 15,000,000 cycles, could be directly measured. For lower frequencies a wave meter was employed. This consisted of a loop 32 centimeters in diameter made of $\frac{1}{8}$ -inch brass rod and a high grade .00035 mfd. condenser in series therewith, resonance being indicated either by the lighting of a low-resistance flash lamp bulb in series with the loop and condenser, or by observing the deflection in the plate-milliammeter needle of the oscillator at the reso-

nance point. This latter method was preferred because of the sharpness of the reaction.

This wave meter was calibrated at the Cruft High Tension Laboratory by comparison with the Cruft Laboratory precision wave-meter for this range of frequencies which, in turn, had been calibrated against a quartz-crystal controlled oscillator.³

Utilization of oscillator output.—In studying the effects of the oscillator output at various frequencies upon laboratory animals it would obviously be inexpedient to make use of any conductive arrangement, as the constants of the oscillating system would thereby be seriously disturbed. In these studies, therefore, the effects of these high frequency currents upon animals were investigated by the use of a tuned circuit which was inductively coupled to the oscillator.

As shown in the circuit diagram (fig. 3) and photograph (Pl. III), this tuned circuit consisted of a single-turn wire loop, having a thermocouple (to measure the current) inserted at its mid-point and a capac-




FIG. 3.—Auxiliary-tuned circuit and thermocouple

ity consisting of a pair of rectangular brass plates 7.4 by 4.3 centimeters, 2 centimeters apart. The separation of the plates was rigidly fixed by means of four $\frac{1}{4}$ -inch hard-rubber posts each 2 centimeters long at each corner of the plates. Since it was determined to study the action of the electrostatic field between the condenser plates, to avoid any conductive transfer of energy the inner surface of each condenser plate was covered with a celluloid sheet 0.004 inch in thickness. The subject for experimentation (a mouse) was placed in a small celluloid box with perforated sides (see Pl. III) and inserted between the condenser plates where it fitted snugly.

The dimensions of the condenser given above were not the result of any calculation, but were determined by the size of the celluloid box which was designed to hold a 20-22 gram mouse comfortably without, at the same time, permitting it to turn and twist too freely. The box consisted of two pieces of stout celluloid for the top and bottom, held

³ See: Piezoelectric Crystal Resonators and Crystal Oscillators Applied to the Precision Calibration of Wave-meters. By George W. Pierce. Proceedings of the American Academy of Arts and Sciences, vol. 50, No. 4.

apart by $\frac{1}{4}$ -inch hard-rubber posts to which they were secured by short screws in holes tapped at each end of the posts. The sides of the box were covered in with strips of stout celluloid perforated with numerous $\frac{1}{4}$ -inch holes for ventilation, glued to the hard-rubber posts by means of cellulose varnish.

While mice of from 19–22 grams in weight fitted snugly in the box, so that they could indulge in but moderate twisting and turning about, still, confinement in the box was apparently not uncomfortable per se, as mice frequently went to sleep when undisturbed. Confinement in the box of several hours' duration was without effect upon mice.

The capacity of the condenser without mouse or mouse holder was found to be 4.1 micromicrofarads. Putting the empty mouse holder in place increased the capacity to 8.1 micromicrofarads. With a 20-gram mouse in the holder the capacity increased to 16.1 micromicrofarads. These measurements were made at 1,000 cycles on the Cruft Laboratory capacity bridge. The capacity of the condenser with mouse holder and mouse was, therefore, about four times greater than the capacity with air alone as the dielectric.

Current measurement.—The amount of oscillating current induced in the auxiliary-tuned circuit was measured by means of a platinum-tellurium thermocouple which was constructed for the writer by Mr. F. H. Drake, of Cruft Laboratory. This consisted of a bit of tellurium supported upon a piece of 22-gauge nickel wire to which it was fused. This in turn was soldered to a short piece of No. 18 copper wire. A piece of 7-mil. platinum wire was rolled to a ribbon approximately 1 mil. in thickness. A piece of this about 1 centimeter long was soldered at one end to a short piece of 18-gauge copper wire, while the other end was made somewhat pointed by means of shears. The pieces of copper wire to which, respectively, the platinum ribbon and the tellurium were attached were mounted vertically, through holes in which they were a tight fit, on a stout piece of bakelite at such distance apart that the pointed end of the platinum ribbon fell naturally about on the center of the tellurium fragment. Upon passing direct current regulated by a variable resistance from a 6-volt storage battery, the platinum ribbon was heated to redness. The tellurium fused at the point of contact and thus became solidly welded to the platinum ribbon. A cap fitted over the thermocouple protected it from the effect of air currents. The circuit diagram of the thermocouple is shown in Figure 3.

It will be seen that the thermocouple is connected across two conductors, two ends of which form part of the auxiliary-tuned circuit, the other ends being connected through choke coils to the plus and minus posts of a Rawson microammeter. This instrument had a resistance of 52 ohms and a full scale reading of 120 microamperes.

The resistance of the thermocouple was found to be 0.3 ohms at 1,000 cycles.

The thermocouple was calibrated on 60-cycle A. C. by noting the scale reading of the microammeter for various current values as determined by the voltage drop across standard resistances measured by a Rawson vacuum thermocouple voltmeter. The full scale read-




FIG. 4.—Resonance curve of auxiliary-tuned circuit with 20-gram mouse. Peak at 10.42 megacycles

ing of 120 on the microammeter corresponded to 0.61 amperes. At 5 on the scale the current was 0.139 amperes.

The calibration of this thermocouple was checked at frequent intervals throughout the study and remained unchanged.

Tuning the auxiliary circuit.—Since the output of the oscillator was applied to the experimental animals, by means of the current

induced in the auxiliary circuit placed in inductive relation to the oscillator, the first step in any experiment was, of course, to tune the coupled circuit to the oscillator frequency so that the current in this circuit should be a maximum. To do this, a mouse immediately after death by gas was placed in the celluloid box and inserted between the condenser plates. A loop was then formed by joining with




FIG. 5.—Resonance curve of auxiliary-tuned circuit with 20-gram mouse. Peak, 28.4 to 27.2 megacycles

two pieces of stiff wire the high frequency terminals of the thermocouple with the corresponding condenser terminals. As a first approximation the length of wire was arbitrarily chosen at a value which experience showed to be about the length desired. Then by starting the oscillator and varying its tuning condenser until maximum current was flowing in the coupled circuit, upon determining the fre-

quency an idea could readily be had of how closely the auxiliary circuit was tuned to the frequency it was desired to study. With this information it was easy to add or to subtract wire from the auxiliary circuit as required, fine adjustment being finally made at the condenser terminals by loosening the set screws which held the wire in place in holes in the arms supporting the condenser and adjusting the length of the inductance by pushing the wires in or pulling them out of the holes. The graphs (Figs. 4 and 5) of the resonance peak of the tuned auxiliary circuit measured at 10,400,000 cycles, and at 28,400,000 cycles, give a good idea of the type of resonance obtained in this circuit.

It will be noted that the circuit tunes more sharply at the lower frequency. In this case the peak of the resonance curve is about 180,000 cycles broad, while, at the higher frequency, the breadth of the peak is about 1,200,000 cycles.


We may, therefore, conclude, since the tuning of the auxiliary circuit is rather broad, that this circuit presents considerable resistance. Hence, only approximate accuracy is necessary in setting the auxiliary circuit to resonance as the current flowing through it, when tuned to somewhere near the oscillator frequency, is ample for experimental purposes.

Effects of Exposure on Laboratory Animals

It will be noted in the auxiliary circuit just described that the experimental animal, insulated in a box of nonconducting material, is placed in the field between the plates of the condenser while the coupled circuit is excited by the oscillator at some particular frequency. Consequently, no free electrons from the external metallic parts of the circuit can enter, nor can they flow out from the body of the experimental animal. The mouse, however, is subjected to a displacement current, in which electrons in the molecules of the body cells will, according to their state of freedom, either pass from molecule to molecule, first in one and then the other direction or, if bound, are stressed in a direction the polarity of which alternates at the oscillator frequency.


The equivalent electric circuit presents at least two possibilities. The first and more probable possibility is that the system may be regarded as two condensers in series connected by a resistance, the first condenser being formed by one plate of the condenser and the body surface of the mouse, in close proximity to it, the intervening celluloid acting as a dielectric. The second condenser is formed, in

similar fashion, by the lower surface of the body, the intervening tissues acting as the connecting resistance thus:



That this equivalent circuit is probable is shown by the fact that we can readily substitute in place of the mouse two metallic plates connected with each other and spaced far enough apart to fit snugly against the top and bottom of the box. It was determined experimentally that two rectangular copper plates each 2.5 by 2 centimeters and spaced 2 centimeters by a connecting copper strip furnished an electrical equivalent for a mouse. If the coupled circuit were tuned to the oscillator frequency with this arrangement in the box, it was found to be approximately in tune if a mouse were substituted. However, the coupled circuit with the substitute in place had naturally a much lower resistance than with a mouse, so that the current registered for a given filament temperature was considerably higher.

Another possibility is that the mouse's body acts in the circuit as a dielectric of poor quality. In this case the electric equivalent would be a condenser shunted by a resistance as shown below.



It has already been mentioned that the insertion of the mouse holder containing a 20-gram mouse between the condenser plates increased the capacity about four times. The effect of the presence of the mouse holder and mouse on the tuning of the circuit is well shown in the following way. When the auxiliary circuit, with the mouse in place, is tuned to the oscillator's frequency, if the mouse holder and mouse are then removed the needle of the microammeter, indicating the current in the tuned circuit, drops to zero. If the oscillator be tuned to resonance with the circuit in this condition, the frequency will be found at a point considerably higher.

In this case, as might be expected, with a perfect dielectric, as air, filling the space between the condenser plates, the resistance of the

circuit is lower, so that the resonance peak is sharper and the current flowing greater for a given oscillator output.

While it may well be that the electrical behavior of the mouse in the circuit partakes of both modes of action just described, still, on the whole, the first state of affairs described seems the more likely.

Effects upon laboratory animals.—Exposure of small laboratory animals, such as white mice, in the manner described, to the rapidly oscillating field between the condenser plates of the auxiliary tuned circuit, causes death usually in a few minutes if the current value be sufficiently great.

The symptoms observed are, in general, as follows: For a short, variable time the mouse is quiescent. This is followed by agitation increasing with the length of exposure. The ears, tail, and paws turn a bright pink which, in many instances, becomes livid or cyanotic as the exposure is prolonged. There is salivation, the nasal secretion is increased; the head and under parts of the body become wet and bedraggled; the paws are covered with beads of moisture. After a variable time convulsions accompanied by convulsive winking take place, dyspnea sets in, and finally respiration ceases. In males there is usually considerable swelling of the genitalia. The heart, however, continues to beat for a little while after respiration ceases.

In the great majority of instances the body of the mouse appears distinctly warm to the hand, and if the rectal temperature be taken immediately after death considerable elevation in the body temperature is found to have taken place, the temperature varying from 42.2° to 43.1° and even 44° C. However, this is not always the case, as death was repeatedly observed to occur in the usual time, and yet the elevation of the body temperature was only moderate, in one instance the rectal temperature not exceeding 39.2° C., a temperature which is not infrequently observed in apparently normal mice. However, in the majority of instances the exposure has caused considerable elevation of the body temperature and we may infer that the primary fatal effect observed consists in raising the body temperature to a degree incompatible with life.

Since the rectal temperature of a normal mouse, according to the temperature of the environment, may be anything from less than 37° to about 39° C., the exposure has brought an increase of the body temperature of anywhere from 5° to 6° C.

In the case of mice, however, which had been killed by exposure to carbon monoxide gas, and then immediately exposed in the coupled circuit, the heating effect was far less pronounced. In dead mice it was found, using the same current and time of exposure necessary to kill living mice at that particular frequency, that the

rise in the body temperature of the dead mouse was of a much lower order, amounting to but 0.7° to 1° C. In many instances there was no gain, and in others a loss was recorded.

This would suggest that the heating effect is different from the diathermic effect observed at the lower frequencies used for therapeutic purposes, as by the use of high-frequency current in diathermic apparatus it is easy to raise the temperature of dead tissues well above the point at which albumen coagulates.

However, at certain frequencies considerable heating effect was observed even in dead mice. Thus, at 6.6 meters, or a frequency of 45,000,000 cycles, 3 dead mice, as the result of exposure lasting 8.5 minutes, showed gains in the rectal temperature of 2.33° 3.6° and 4.4° C., respectively.

Sequelæ of exposure to sublethal doses.—Apart from the acute symptoms previously mentioned exposure to these high-frequency currents may cause destruction of tissue. After sublethal exposures, in many instances, small hemorrhagic areas may be observed along the course of blood vessels of the ears. In the course of 48 hours the ears become necrotic and drop off. The tail also often presents numerous ecchymotic areas. It may subsequently become affected with dry gangrene and drop off. In other instances areas of alopecia develop, particularly in the supra-orbital region or at the tip of the snout. In one instance a panophthalmitis developed with subsequent loss of vision.

Effects of change in frequency.—In preliminary observations it soon became evident that the effects of exposure to different frequencies was not the same, a current of constant value proving more lethal at some frequencies than at others. Furthermore, the change in lethal effect did not appear to bear any simple relation to change in frequency, as one would be led to expect, so it was determined to study the changes, if any, in the lethality of a constant current as the frequency changed. Since part, at least, of the effect of the exposure was due to increase in the body temperature which, in turn, might be modified by changing external temperature conditions, it was evident that, to avoid serious error, the observations must all be made under substantially constant meteorological conditions.

The standard conditions of temperature and humidity at which to conduct the experiments were arbitrarily fixed at 24° C. and 40 to 42 per cent relative humidity, as these represented a fair average of the atmospheric conditions at which preliminary observations were made.

To secure these conditions, the apparatus depicted in Figure 6 was set up. As will be seen from the diagram, it consisted of the following parts:

The condenser, in the field of which the mouse was exposed, was mounted in a wooden box 22 by 10 by 9.5 centimeters outside dimen-




FIG. 8.—Constant temperature and humidity apparatus. A, Humidifier; B, Hot-water bath; C, Box containing condenser; D, Condenser; E, Mouse holder

sions. The front and back sides of the box, provided with celluloid windows for observation, were removable and were held in place by stout rubber bands. To secure a close fit, the inner side of each was fitted with a rubber gasket 1 centimeter wide which bore against the edges of the box.

Air taken from the laboratory compressed-air supply was first cooled and partly saturated by bubbling through water in a large wash bottle, which itself was cooled to approximately 10° C. by immersion in water of that temperature. The air issuing from the wash bottle was piped to a copper coil inside a large container packed in sawdust and filled with water which was kept at a constant temperature of 10° C. From the copper coil the air issued in a stream of fine bubbles, from a bubbler consisting of a small, weighted tin can, the open end of which was covered with 8 or 10 thicknesses of butter cloth. The air, cooled to 10° C., and thus practically fully saturated with moisture at that temperature, was piped through a jacketed copper pipe to the box in which the condenser was mounted, entering at the bottom through a glass tube in a perforated rubber stopper tightly fitted in a hole in the side of the box.

The ventilating current, rising through the box, issued through a similar glass tube at the top of the opposite side. A ventilation of about 8 to 9 liters per minute was maintained through the box. A centigrade thermometer inserted through a perforated rubber stopper in the top of the box indicated the interior temperature.

Since the amount of moisture which fully saturates air at 10° C. will leave it only 42.1 per cent saturated at 24° C., it is obvious that, to secure the standard atmospheric conditions determined upon, all that is necessary is to see that the temperature of the box is maintained at 24° C., or as close to that figure as is practicable.

In order to provide means of heating the air when necessary, a coil of copper pipe in a hot-water bath was provided in the vicinity of the box, through which air could be by-passed, before reaching the box, by means of two stopcocks, as shown in the figure. In this manner, whenever, owing to external conditions, the temperature of the box tended to fall below 24° C., the incoming air could be led first through the copper coil in the hot-water bath and thus heated to the necessary extent. Since, when heated, no moisture would be lost, nor could it gain any, upon cooling to the temperature of the box, the relative humidity would comply with the conditions.

With the apparatus just described a large number of observations were made at a constant current and at frequencies varying from 135,000,000 to 8,330,000 cycles per second. The constant current employed caused a deflection of 30 divisions on the scale of the microammeter, corresponding to a current of 338 milliamperes flowing in the auxiliary tuned circuit.

This value of current was chosen because preliminary observations had shown that, at lethal frequencies, this current value always caused death in 10 minutes or less. In the great majority of instances ten mice weighing between 19 and 22 grams were used successively at each frequency and sometimes more were used.

The difference in frequency employed was in the neighborhood of 10 per cent. This accounts for the fact that many of the frequencies and wave lengths given in Table 1 are not integers. In that table are set forth the frequencies at which observations were made, the number of observations, the average time elapsing before death, the maximum time, the minimum time, and the usual time.

TABLE 1

| Fre- quency times 10 ⁶ | Wave length | Number of obser- vations | Average length of survival | Mini- mum length of survival | Maxi- mum length of survival | Usual length of survival |
|---|----------------|--------------------------------|----------------------------------|---------------------------------------|---------------------------------------|--------------------------------|
| | <i>Meters</i> | | <i>Minutes</i> | <i>Minutes</i> | <i>Minutes</i> | <i>Minutes</i> |
| 135 | 2.22 | 8 | 26.4 | 16.5 | (¹) | 21-25 |
| 121.5 | 2.47 | 10 | 20.1 | 13.5 | 26.5 | 20-22.5 |
| 110 | 2.73 | 10 | 15.9 | 11.5 | 20.0 | 17 |
| 99 | 3.03 | 11 | 15.6 | 9.25 | 23.0 | 14.5-16 |
| 90 | 3.34 | 12 | 10.9 | 8.0 | 17.5 | 9-12.5 |
| 81 | 3.7 | 9 | 9.9 | 7.0 | 16.5 | 8-10.5 |
| 73 | 4.11 | 10 | 9.5 | 6.5 | 15.5 | 9-9.5 |
| 66 | 4.54 | 10 | 7.7 | 6.0 | 9.0 | 7-8 |
| 60 | 5.0 | 9 | 8.0 | 5.5 | 9.5 | 7-9 |
| 55 | 5.44 | 10 | 9.25 | 5.0 | 12.0 | 9-10 |
| 50 | 6.0 | 6 | 9.37 | 7.0 | 11.0 | 8.5-10 |
| 45 | 6.66 | 14 | 7.8 | 4.5 | 10.5 | 7-8.5 |
| 42.8 | 7.0 | 5 | 8.1 | 6.0 | 11.5 | 7-8 |
| 40.1 | 7.25 | 10 | 7.6 | 6.0 | 9.5 | 7 |
| 35 | 8.51 | 16 | 9.1 | 6.5 | 14.5 | 7.5-9.5 |
| 30 | 10.0 | 10 | 8.5 | 7.0 | 10.5 | 7.5-8.5 |
| 27 | 11.11 | 10 | 6.65 | 5.0 | 8.0 | 6-7 |
| 24.3 | 12.3 | 10 | 10.52 | 8.5 | 14.0 | 9.5-10.25 |
| 22.22 | 13.5 | 13 | 7.8 | 5.0 | 12.0 | 6.5-7.5 |
| 20 | 15.0 | 12 | 8.27 | 6.0 | 11.5 | 7-8.5 |
| 18 | 16.6 | 15 | 6.95 | 4.5 | 9.5 | 5.5-6.5 |
| 16.2 | 18.5 | 11 | 7.1 | 5.5 | 11.5 | 6-7.5 |
| 14.6 | 20.55 | 9 | 11.3 | 7.75 | 15.0 | 10.5-12 |
| 13.7 | 21.9 | 10 | 10.6 | 7.5 | 16.5 | 10-12 |
| 12.33 | 24.4 | 10 | 14.25 | 9.0 | 19.5 | 12-14.5 |
| 11.1 | 27.0 | 11 | 14.5 | 8.0 | 19.5 | 12-13.5 |
| 9.93 | 30.2 | 10 | 9.65 | 6.5 | 14.0 | 8-9 |
| 9.2 | 32.5 | 9 | 11.5 | 7.5 | 17.0 | 10-11.5 |
| 9.0 | 33.3 | 10 | 23.5 | 15.5 | ² 35.0 | 22-30 |
| 8.5 | 35.3 | 11 | 17.2 | 11.0 | ³ 30.0 | 16-18 |
| 8.33 | 36.0 | 11 | 17.7 | 11.5 | 32.0 | 16-19 |

¹ 1 mouse survived 43 minutes unaffected.

² 3 mice survived 35 minutes unaffected.

³ 1 mouse survived 30 minutes unaffected.

The average time of survival plotted against frequency is graphically shown in the chart which accompanies the table.

It will be seen from the graph that, starting at a frequency of 135,000,000 cycles per second, with a constant current maintained in the auxiliary circuit of 338 milliamperes, the lethality increases until at 66,000,000 cycles the mean time of survival has been reduced from an average of 26.8 minutes to one of 7.7 minutes, a reduction of 71.3 per cent. From this point, with a fluctuation of a minute or so in the mean time of survival, the lethality of the various frequencies




Chart showing the average time of survival of mice subjected to currents of various frequencies

observed maintains itself at more or less the same value over a frequency band of considerable width, i. e. from 66,000,000 to 27,000,000 cycles, at which point the low average period of survival of 6.65 minutes is recorded, or about 75 per cent less than the average time of survival at 135,000,000 cycles. At 24.3 million cycles the current becomes less lethal by about 40 per cent but at 18,000,000 cycles the current has about the same degree of lethality as at 27,000,000 cycles.

From 18,000,000 cycles the lethality on the whole diminishes, till at 11.1 million cycles the average time of survival increases to 14.5 minutes, 105 per cent longer than at 18,000,000 cycles. At a frequency of 10,000,000 cycles a recrudescence in lethality is observed, the mean time of survival falling to 9.65 minutes. At a frequency of 9,000,000 cycles, however, the lethality is comparatively low, the average time of survival being 23.5 minutes, nearly the same as it was at 135,000,000 cycles. At this frequency three mice survived exposures lasting 30 to 35 minutes apparently unharmed except for slight elevations of the body temperature. At 8.33 million cycles, beyond which point the observations were not carried, the lethality of the exposure seems on the increase, as well as at 8.5 million cycles, as the average time of survival of 11 mice was 17.2 and 17.7 minutes, respectively.

Discussion

From the configuration of the curve it is evident that the results just described are unexpected. Starting with a low lethal effect at the highest frequencies studied, a gradual increase in lethality is observed, followed by maintenance of fatality of the same order with successive small maxima and minima until at 18,000,000 cycles a peak is reached, followed by an abrupt decline, separated from the low point at 9,000,000 cycles by another maximum.

This result differs from what is usually observed in physical phenomena of this character. On the whole one would expect to find either that the lethality of a constant current was independent of frequency, or if dependent would show some simple relation thereto, e. g., be either inversely or directly proportional. That the lethality should be at first inversely and subsequently directly proportional to frequency is puzzling, and certainly not susceptible of any simple explanation.

At first one is tempted to assume that the results of the observation are apparent only. For instance, it might be urged that the apparent low lethality of the higher frequencies might be due to the "skin" effect of these frequencies upon the thermocouple.

As is well known, the high frequency resistance of conductors increases with the frequency, because, as this becomes greater, the current is confined to a surface pellicle of ever-diminishing thickness.

This so-called "skin effect" would occasion an increase with frequency in the apparent resistance of the thermocouple. Hence there would be a greater heating effect at the very high frequencies, and, as a result, a higher e. m. f. between the hot and cold junctions of the thermocouple, which would be reflected by a higher reading on the microammeter scale than would be justified by the actual oscillatory current flowing in the circuit.

That there may be some such effect at the highest frequencies is not denied; but how explain on these grounds the pronounced recession in lethality at the lower frequencies, where the "skin effect" is admittedly much less. Thus, we note that the lethality at 18.1 million cycles is about the same as at 24.3 million cycles, but very much greater (nearly 240 per cent) than at 9,000,000 cycles. Yet the skin effect, if important, should be more noticeable at 18,000,000 than at 9,000,000 cycles.

In view of the circumstance that, generally speaking, exposure to the oscillating current has very much less heating effect on dead than on living tissues, it may be suggested that, here, we are dealing with some action other than the direct heating effect observed at ordinary frequencies with diathermic apparatus.

In this case the heating effect is proportional to the product of the resistance and the square of the current. Under the conditions of this study it is possible that the heat generated is caused by dielectric loss or hysteresis.

Again there is the possibility that the action is indirect rather than direct, the high-frequency current increasing the basal rate of metabolism, or perhaps causing heat retention through its influence upon thermogenic centers.

That the effects of exposure to the high-frequency current is capable of directly damaging tissue is shown by the sequelae already described, which are observed in mice subjected to a sublethal exposure.

As for the differential action of the different frequencies, which is the most striking of the effects observed, nothing except of a speculative nature can be advanced by way of explanation. For instance, it might be thought possible that at certain frequencies harmonics generated by the oscillator are in resonance with the natural period of organic molecules, particularly of animal proteids. These are well known to possess a very high molecular weight except for certain colloidal aggregates, perhaps the highest known molecular weights. Consequently, the natural period of resonance would be correspondingly low.

Nevertheless, in spite of the great size of the body molecules, the natural period of these, though relatively low compared with the molecules of simpler structure, would be sufficiently high to correspond to a wave length of from 0.5 to 3 or 4 millimeters, and hence to a fre-

quency of six times 10^{10} to seven times 10^9 . The generation of definite harmonics of such a high order on the part of the oscillator seems altogether unlikely.

I am, however, indebted to Prof. George W. Pierce for the suggestion that the differential effects due to frequency may be caused by a mode of electromechanical vibration of the tissue cells, and for the accompanying reference to literature on the subject. Mechanical oscillations in solid elastic spheres have been given thorough mathematical treatment by Lamb.⁴ A number of modes of vibration are analyzed and certain constants corresponding therewith are given. Some of the values of these constants (here designated " K ") as given by Lamb, are: $K=0.66, 1.89, 2.93, 3.94, 4.96, 5.97$, etc., the lowest value corresponding to the mode of vibration having the longest period. Since a solid elastic sphere is capable of vibrating in a number of different successive modes, each of which has a frequency greater than that preceding it, such a sphere is capable of responding to impressed oscillations of different frequencies.

The frequency, F , at which the sphere vibrates in a particular mode is given in simplest form by the equation—

$$F = K \frac{V}{D} \quad (1)$$

where F = Frequency of the particular mode of vibration.

K = The corresponding constant given by Lamb.

V = Velocity of the propagation of sound in the material composing the sphere.

D = Diameter of the sphere.

From this equation it follows directly that the diameter of a sphere vibrating in a particular mode at frequency, F , is given by the equation—

$$D = K \frac{V}{F} \quad (2)$$

In this case let us assume that the velocity of the propagation of sound through tissue cells is approximately that of its propagation through water, i. e., 1.45×10^5 centimeters per second. For the value of K let us take 0.66, which corresponds to the mode of vibration having the longest period and thus to the diameter of the smallest sphere capable of vibrating in some mode at that particular frequency. For F let us assume a value $F = 40 \times 10^9$, a frequency somewhere in the middle of the band of lethal frequencies observed. Substituting these values in equation (2) we obtain for D , diameter of the sphere, 0.0024 centimeters or 24 microns. This diameter, while great for body cells, is, nevertheless, encountered in a number

⁴ Lamb, H.: On the Vibrations of an Elastic Sphere. Proceedings of the London Mathematical Society Series, 1, 1882, vol. 13, p. 189.

of cells in the nervous system. The diameter of the smallest sphere capable of oscillating at the highest frequencies worked with would be in the neighborhood of 7 microns, while at the lowest lethal frequency (lowest frequency of high lethality 10×10^6), this diameter would be 96 microns. This is greater than the diameter of any body cells except, perhaps, some of the giant pyramidal cells of the nervous system. However, in this instance one might venture the speculation that second or third order harmonics of the oscillator might, at the lower frequencies, possess sufficient energy to induce some form of electromechanical resonance in the body cells, although the fundamental frequency is too low to resonate with any except the sufficiently large body cells just mentioned.

As to the way in which electrical oscillations could induce mechanical vibrations in body cells, speculations are again in order. One possibility, at least, is that cell membranes act as dielectrics at the boundaries of cells. This would cause the contiguous surfaces of cells to act as condensers resulting in stresses of alternating polarity at the impressed frequency. A reinforcement of the effect resulting in mechanical vibrations would reasonably be expected when the frequency of the applied stresses approaches the natural period for mechanical vibrations of the cell. It would not, however, be expected that this frequency is critical, as such a system must, of necessity, be highly damped. So, if responsive, the response would merely be greater for a range of frequencies in the neighborhood of the frequency of mechanical vibrations.

Conclusions

On the basis of the data gathered in the foregoing experiments, the following conclusions are permitted:

1. When small laboratory animals are placed in a box of insulating material and subjected to the action of a high frequency oscillating current in the field of a condenser resonating a tuned circuit, severe symptoms are caused which may result in death if the exposure is prolonged. Part at least, of the symptoms is due to heat retention.

2. This effect is most marked in a certain band of frequencies extending from $F=66 \times 10^6$ cycles to $F=18.3 \times 10^6$ cycles, the effect diminishing in one direction from a band extending from $F=66 \times 10^6$ to $F=135 \times 10^6$ and in the other from $F=18.3 \times 10^6$ to $F=9 \times 10^6$.

3. There is, consequently, at constant current, under the conditions of the experiments, a differential action with respect to frequency, the lethality of a constant current being in one region of the spectrum inversely and in another portion directly proportional to frequency.

4. In the band of frequencies studied, successive maxima and minima with respect to lethality occur which are most pronounced as the lower frequencies are approached.

Finally it may be remarked that here we are dealing with a band in the spectrum of radiant energy which as yet has been little studied in its effects on living cells. Since frequency is the sole differentiating characteristic in the whole band of radiant energy it is perhaps to be expected to find that in electromagnetic waves, frequency is a determining factor in their mode of action upon living organisms. It is thought that this is a field which will well repay further study.

PUBLIC HEALTH ENGINEERING ABSTRACTS

Report on the First Results of Laboratory Work on Malaria in England. S. P. James. Publication by the Health Section of the League of Nations, Geneva, 1926. (Abstract by L. D. Fricks.)

While this is a report of observations made on artificially infected mosquitoes in the laboratory, it contains valuable suggestions to those who are engaged in malaria control. These observations were made in connection with the malarial treatment of general paralysis. Two thousand six hundred and thirty-eight *Anopheles maculipennis* were employed, 532 of which became infected and were used to bite 145 persons, 109 of whom developed malaria. Colonel James observes that it is no easy matter to infect mosquitoes with malaria even under the most favorable laboratory conditions. The conditions most favorable to the development of the *Plasmodia* in the mosquito causes high mortality among the mosquitoes themselves. To produce infective mosquitoes it is necessary that they be kept, after feeding on a malaria carrier, at a temperature of 24° C. in an atmosphere saturated with moisture, and the mosquitoes must be given frequent feedings—daily or every other day. It would seem that the required conditions for producing naturally infected mosquitoes are only rarely met. In addition, very few malaria patients are good infectors of *Anopheles*. The extreme delicacy of these requirements indicate to the author that malaria should be dealt with in the houses of the people rather than in the environment, and that a waste of effort is involved in measures directed toward general mosquito destruction.

(Abstractor's note: LePrince pointed out the importance of destroying engorged *Anopheles* in houses as a malaria-control measure in the Panama Canal Zone many years ago.)

Problems in Malaria Control. W. E. Deeks. Fourteenth Annual Report United Fruit Company, 1925, pp. 170-186. (Abstract by L. D. Fricks.)

Doctor Deeks states that malaria is responsible for 40 per cent of the sickness on the plantations of the United Fruit Company, and it is therefore the most important single factor in lowering the efficiency of labor. A description is given of certain United Fruit Company plantations and the extent of the malaria problem thereon indicated. Among the conditions which influence the incidence of

malaria on these plantations are listed the following: Rainfall; location of camps; screening of quarters; and impaired physical condition frequently due to complicating diseases. Measures of malaria control recommended are: Mosquito control for a radius of two miles around the settlements; careful selection of new camp sites; stabilizing of population; and screening of houses and cure of carriers.

AMERICAN PUBLIC HEALTH ASSOCIATION TO MEET IN BUFFALO, OCTOBER 11-14, 1926

What is new in public health? How are communities, urban and rural, coping with the public health problems that confront administrators, public health officers, nurses, and inspectors? These questions will be answered, and problems that have been the subject of laboratory research will be discussed by specialists at the fifty-fifth annual meeting of the American Public Health Association to be held in Buffalo, N. Y., October 11-14, 1926. This association is the professional society of the public health workers of North America.

It is expected that this year's meeting will be the largest and the most interesting of all the annual gatherings of the association, and this expectation is fully justified on the basis of the announcements contained in the preliminary program recently issued. The sessions are arranged in four sections—general sessions; special sessions, which have been developed around subjects of timely and lively interest; sessions of the nine scientific sections of the association; and the special program arranged by the New York State Conference of Health Officers and Public Health Nurses, which the State health commissioner, Dr. Matthias Nicoll, has called to meet in conjunction with the American Public Health Association.

The meeting will begin on Monday morning, October 11, and will end with a special dinner session on Thursday evening, October 14. The headquarters of the association will be in the Hotel Statler, where all sessions will be held. The preliminary program and other information regarding the meeting may be had by addressing Homer N. Calver, Executive Secretary, American Public Health Association, 370 Seventh Avenue, New York City.

AMERICAN DIETETIC ASSOCIATION MEETING AT ATLANTIC CITY

The American Dietetic Association will hold its ninth annual meeting at Atlantic City, October 11, 12, and 13, 1926. The program will include papers and discussions on both scientific and administrative matters.

Dr. J. J. R. McLeod, of the University of Toronto, will address the convention on the advances made in physiology, and Dr. Julius Stieglitz, of the University of Chicago, will speak on the recent advances made in chemistry.

Arrangements have been made for some interesting exhibits, both commercial and noncommercial.

Dr. Ruth Wheeler, professor of physiology at Vassar, is the president of the association.

DEATHS DURING WEEK ENDED AUGUST 28, 1926

Summary of information received by telegraph from industrial insurance companies for week ended August 28, 1926, and corresponding week of 1925. (From the Weekly Health Index, September 1, 1926, issued by the Bureau of the Census, Department of Commerce)

| | Week ended Aug. 28, 1926 | Corresponding week, 1925 |
|--|-----------------------------|-----------------------------|
| Policies in force..... | 64, 869, 549 | 60, 879, 605 |
| Number of death claims..... | 10, 175 | 10, 590 |
| Death claims per 1,000 policies in force, annual rate..... | 8. 2 | 9. 1 |

Deaths from all causes in certain large cities of the United States during the week ended August 28, 1926, infant mortality, annual death rate, and comparison with corresponding week of 1925. (From the Weekly Health Index, September 1, 1926, issued by the Bureau of the Census, Department of Commerce)

| City | Week ended Aug. 28, 1926 | | Annual death rate per 1,000 corresponding week, 1925 | Deaths under 1 year | | Infant mortality rate, week ended Aug. 28, 1926 ¹ |
|------------------------|--------------------------|-------------------------|--|--------------------------|--------------------------|--|
| | Total deaths | Death rate ¹ | | Week ended Aug. 28, 1926 | Corresponding week, 1925 | |
| Total (65 cities)..... | 5, 625 | 10. 2 | 10. 5 | 760 | 901 | ¹ 61 |
| Akron..... | 27 | ----- | 13. 3 | 8 | 11 | 85 |
| Albany..... | 19 | 8. 3 | ----- | 1 | 0 | 21 |
| Atlanta..... | 63 | ----- | ----- | 13 | 7 | ----- |
| White..... | 26 | ----- | ----- | 4 | ----- | ----- |
| Colored..... | 37 | (⁴) | ----- | 9 | ----- | ----- |
| Baltimore..... | 167 | 10. 8 | 10. 7 | 25 | 22 | 73 |
| White..... | 125 | ----- | ----- | 17 | ----- | 61 |
| Colored..... | 42 | (³) | ----- | 8 | ----- | 130 |
| Birmingham..... | 56 | 13. 8 | 14. 7 | 13 | 6 | ----- |
| White..... | 34 | ----- | ----- | 7 | ----- | ----- |
| Colored..... | 22 | (⁵) | ----- | 6 | ----- | ----- |
| Boston..... | 186 | 12. 3 | 11. 5 | 45 | 23 | 127 |
| Bridgeport..... | 28 | ----- | ----- | 3 | 2 | 61 |
| Buffalo..... | 119 | 11. 4 | 10. 7 | 8 | 24 | 33 |
| Cambridge..... | 18 | 7. 7 | 7. 8 | 3 | 3 | 50 |
| Camden..... | 23 | 9. 2 | 8. 1 | 7 | 4 | 118 |
| Canton..... | 21 | 10. 0 | 3. 9 | 4 | 0 | 89 |
| Chicago..... | 533 | 9. 1 | 9. 3 | 61 | 81 | 54 |
| Cincinnati..... | 129 | 16. 4 | 14. 0 | 15 | 14 | 93 |
| Cleveland..... | 166 | 9. 0 | 9. 6 | 21 | 36 | 54 |
| Columbus..... | 64 | 11. 7 | 11. 9 | 10 | 12 | 92 |
| Dallas..... | 39 | 10. 2 | 10. 5 | 8 | 7 | ----- |
| White..... | 28 | ----- | ----- | 8 | ----- | ----- |
| Colored..... | 11 | (⁵) | ----- | 0 | ----- | ----- |
| Dayton..... | 38 | 11. 2 | 9. 6 | 0 | 2 | 0 |
| Denver..... | 62 | 11. 3 | 16. 0 | 8 | 16 | ----- |
| Des Moines..... | 16 | 5. 7 | 8. 5 | 2 | 3 | 33 |
| Detroit..... | 212 | 8. 6 | 10. 4 | 39 | 66 | 63 |
| Duluth..... | 21 | 9. 7 | 9. 0 | 2 | 3 | 47 |
| El Paso..... | 23 | 11. 0 | 11. 9 | 3 | 5 | ----- |
| Erie..... | 21 | ----- | ----- | 2 | 1 | 38 |
| Fall River..... | 24 | 9. 6 | 9. 7 | 2 | 5 | 29 |
| Flint..... | 17 | 6. 5 | 7. 2 | 5 | 5 | 83 |
| Fort Worth..... | 25 | 8. 2 | 6. 2 | 5 | 1 | ----- |
| White..... | 22 | ----- | ----- | 5 | ----- | ----- |
| Colored..... | 3 | (⁴) | ----- | 0 | ----- | ----- |
| Grand Rapids..... | 19 | 6. 4 | 11. 2 | 0 | 10 | 0 |
| Houston..... | 37 | ----- | ----- | 4 | 4 | ----- |
| White..... | 28 | ----- | ----- | 2 | ----- | ----- |
| Colored..... | 9 | (⁵) | ----- | 2 | ----- | ----- |
| Indianapolis..... | 92 | 13. 1 | 12. 1 | 12 | 10 | 88 |
| White..... | 73 | ----- | ----- | 8 | ----- | 68 |
| Colored..... | 19 | (⁵) | ----- | 4 | ----- | 220 |

See footnotes at end of table.

Deaths from all causes in certain large cities of the United States during the week ended August 28, 1926, infant mortality, annual death rate, and comparison with corresponding week of 1925—Continued

| City | Week ended Aug. 28, 1926 | | Annual death rate per 1,000 corresponding week, 1925 | Deaths under 1 year | | Infant mortality rate, week ended Aug. 28, 1926 ¹ |
|------------------------|--------------------------|-------------------------|--|--------------------------|--------------------------|--|
| | Total deaths | Death rate ¹ | | Week ended Aug. 28, 1926 | Corresponding week, 1925 | |
| Jersey City..... | 40 | 6.6 | 10.4 | 4 | 8 | 28 |
| Kansas City, Kans..... | 21 | 9.4 | 12.1 | 0 | 2 | 0 |
| White..... | 17 | | | 0 | | 0 |
| Colored..... | 4 | (²) | | 0 | | 0 |
| Kansas City, Mo..... | 85 | 11.8 | 11.4 | 13 | 11 | |
| Los Angeles..... | 203 | | | 15 | 22 | 42 |
| Louisville..... | 74 | 12.4 | 10.7 | 9 | 8 | 78 |
| White..... | 54 | | | 7 | | 70 |
| Colored..... | 20 | (²) | | 2 | | 125 |
| Lowell..... | 18 | | | 2 | 6 | 37 |
| Lynn..... | 15 | 7.5 | 6.1 | 2 | 3 | 50 |
| Memphis..... | 61 | 18.0 | 16.7 | 9 | 9 | |
| White..... | 29 | | | 5 | | |
| Colored..... | 32 | (²) | | 4 | | |
| Milwaukee..... | 74 | 7.5 | 8.8 | 16 | 7 | 74 |
| Minneapolis..... | 92 | 11.1 | 9.7 | 9 | 9 | 50 |
| Nashville..... | 55 | 20.9 | 14.9 | 8 | 3 | |
| New Bedford..... | 31 | | | 5 | 5 | 87 |
| New Haven..... | 53 | 15.2 | 11.4 | 5 | 8 | 68 |
| New Orleans..... | 147 | 18.3 | 16.6 | 17 | 21 | |
| White..... | 70 | | | 5 | | |
| Colored..... | 71 | (²) | | 12 | | |
| New York..... | 1,036 | 9.1 | 9.7 | 126 | 154 | 51 |
| Bronx Borough..... | 123 | 7.1 | 6.9 | 10 | 8 | 33 |
| Brooklyn Borough..... | 341 | 7.9 | 8.6 | 54 | 57 | 55 |
| Manhattan Borough..... | 425 | 11.8 | 12.9 | 45 | 73 | 50 |
| Queens Borough..... | 103 | 7.0 | 7.2 | 16 | 15 | 73 |
| Richmond Borough..... | 44 | 16.0 | 10.9 | 1 | 1 | 18 |
| Newark, N. J..... | 101 | 11.5 | 10.7 | 25 | 15 | 120 |
| Norfolk..... | 32 | 9.6 | 8.0 | 2 | 3 | 37 |
| White..... | 8 | | | 0 | | 0 |
| Colored..... | 24 | (²) | | 2 | | 90 |
| Oakland..... | 43 | 8.6 | 8.8 | 7 | 6 | 81 |
| Oklahoma City..... | 18 | | | 3 | 0 | |
| Omaha..... | 53 | 12.8 | 13.1 | 6 | 5 | 63 |
| Paterson..... | 18 | 6.6 | 12.1 | 3 | 2 | 52 |
| Philadelphia..... | 354 | 9.2 | 10.6 | 40 | 77 | 53 |
| Pittsburgh..... | 147 | 12.0 | 13.3 | 23 | 33 | 76 |
| Portland, Oreg..... | 52 | | | 2 | 3 | 20 |
| Providence..... | 53 | 10.0 | 8.8 | 7 | 8 | 58 |
| Richmond..... | 43 | 11.9 | 11.5 | 12 | 7 | 151 |
| White..... | 24 | | | 4 | | 78 |
| Colored..... | 19 | (²) | | 8 | | 280 |
| Rochester..... | 64 | 10.4 | 11.9 | 5 | 12 | 40 |
| St. Louis..... | 138 | 9.9 | 10.7 | 21 | 27 | |
| St. Paul..... | 41 | 8.6 | 12.3 | 1 | 3 | 9 |
| Salt Lake City..... | 29 | 11.4 | 8.8 | 0 | 0 | 0 |
| San Antonio..... | 45 | 11.4 | 12.4 | 6 | 7 | |
| San Diego..... | 29 | 13.7 | 10.3 | 1 | 1 | 21 |
| San Francisco..... | 116 | 10.7 | 11.0 | 8 | 11 | 48 |
| Schenectady..... | 24 | 13.5 | 12.4 | 6 | 8 | 173 |
| Seattle..... | 61 | | | 4 | 5 | 37 |
| Somerville..... | 19 | 9.9 | 8.4 | 6 | 2 | 156 |
| Spokane..... | 17 | 10.1 | 10.5 | 4 | 1 | 94 |
| Springfield, Mass..... | 28 | 10.1 | 9.2 | 3 | 2 | 43 |
| Syracuse..... | 43 | 12.2 | 10.3 | 3 | 7 | 38 |
| Toledo..... | 52 | 9.2 | 10.5 | 7 | 8 | 68 |
| Trenton..... | 31 | 12.1 | 13.0 | 4 | 8 | 67 |
| Utica..... | 25 | 12.7 | 10.3 | 2 | 3 | 44 |
| Washington, D. C..... | 109 | 10.8 | 12.3 | 15 | 22 | 85 |
| White..... | 72 | | | 7 | | 58 |
| Colored..... | 37 | (²) | | 8 | | 146 |
| Waterbury..... | 23 | | | 5 | 4 | 107 |
| Wilmington, Del..... | 18 | 7.6 | 8.5 | 4 | 5 | 94 |
| Worcester..... | 37 | 10.0 | 12.6 | 9 | 7 | 104 |
| Yonkers..... | 15 | 6.7 | 11.5 | 2 | 5 | 45 |
| Youngstown..... | 39 | 12.3 | 7.8 | 11 | 5 | 140 |

¹ Annual rate per 1,000 population.

² Deaths under 1 year per 1,000 births. Cities left blank are not in the registration area for births.

³ Data for 63 cities.

⁴ Deaths for week ended Friday, Aug. 27, 1926.

⁵ In the cities for which deaths are shown by color, the colored population in 1920 constituted the following percentages of the total population: Atlanta 31, Baltimore 15, Birmingham 39, Dallas 15, Fort Worth 14, Houston 25, Indianapolis 11, Kansas City, Kans., 14, Louisville 17, Memphis 38, New Orleans 26, Norfolk 38, Richmond 32, and Washington, D. C., 25.

PREVALENCE OF DISEASE

No health department, State or local, can effectively prevent or control disease without knowledge of when, where, and under what conditions cases are occurring

UNITED STATES

CURRENT WEEKLY STATE REPORTS

These reports are preliminary, and the figures are subject to change when later returns are received by the State health officers

Reports for Week Ended September 4, 1926

| ALABAMA | | CALIFORNIA | |
|-------------------------------|-------|--|-------|
| | Cases | | Cases |
| Cerebrospinal meningitis..... | 1 | Cerebrospinal meningitis: | |
| Chicken pox..... | 4 | Long Beach..... | 1 |
| Diphtheria..... | 18 | San Diego County..... | 1 |
| Influenza..... | 8 | Chicken pox..... | 29 |
| Malaria..... | 112 | Diphtheria..... | 87 |
| Measles..... | 20 | Influenza..... | 6 |
| Mumps..... | 7 | Measles..... | 141 |
| Ophthalmia neonatorum..... | 1 | Mumps..... | 60 |
| Pellagra..... | 11 | Poliomyelitis: | |
| Pneumonia..... | 25 | Long Beach..... | 1 |
| Scarlet fever..... | 12 | Paso Robles..... | 1 |
| Smallpox..... | 7 | Redwood City..... | 1 |
| Tuberculosis..... | 56 | San Diego..... | 1 |
| Typhoid fever..... | 103 | Rabies (human)—Los Angeles County..... | 1 |
| Whooping cough..... | 17 | Scarlet fever..... | 63 |
| | | Smallpox..... | 7 |
| ARIZONA | | Tuberculosis..... | 163 |
| Diphtheria..... | 7 | Typhoid fever..... | 20 |
| Malaria..... | 1 | Whooping cough..... | 40 |
| Measles..... | 2 | | |
| Tuberculosis..... | 16 | COLORADO | |
| Typhoid fever..... | 1 | Chicken pox..... | 10 |
| ARKANSAS | | Diphtheria..... | 13 |
| Chicken pox..... | 12 | Influenza..... | 4 |
| Diphtheria..... | 3 | Mumps..... | 2 |
| Hookworm disease..... | 2 | Pneumonia..... | 1 |
| Influenza..... | 13 | Scarlet fever..... | 4 |
| Malaria..... | 198 | Tuberculosis..... | 54 |
| Measles..... | 3 | Typhoid fever..... | 15 |
| Mumps..... | 8 | Vincent's angina..... | 2 |
| Paratyphoid fever..... | 2 | Whooping cough..... | 27 |
| Pellagra..... | 11 | | |
| Poliomyelitis..... | 1 | CONNECTICUT | |
| Scarlet fever..... | 1 | Cerebrospinal meningitis..... | 1 |
| Smallpox..... | 5 | Chicken pox..... | 2 |
| Trachoma..... | 2 | Diphtheria..... | 9 |
| Tuberculosis..... | 15 | German measles..... | 1 |
| Typhoid fever..... | 92 | Influenza..... | 1 |
| Whooping cough..... | 19 | Malaria..... | 1 |

(1967)

CONNECTICUT—continued

| | Cases |
|-------------------------------|-------|
| Measles..... | 9 |
| Mumps..... | 1 |
| Paratyphoid fever..... | 1 |
| Pneumonia (broncho)..... | 7 |
| Pneumonia (lobar)..... | 6 |
| Polioimyelitis..... | 1 |
| Scarlet fever..... | 13 |
| Septic sore throat..... | 18 |
| Tuberculosis (all forms)..... | 35 |
| Typhoid fever..... | 8 |
| Whooping cough..... | 36 |

DELAWARE

| | |
|---------------------|---|
| Diphtheria..... | 1 |
| Malaria..... | 1 |
| Polioimyelitis..... | 1 |
| Scarlet fever..... | 4 |
| Tuberculosis..... | 2 |
| Typhoid fever..... | 2 |

FLORIDA

| | |
|---------------------|----|
| Diphtheria..... | 10 |
| Influenza..... | 2 |
| Malaria..... | 6 |
| Measles..... | 5 |
| Mumps..... | 6 |
| Scarlet fever..... | 4 |
| Smallpox..... | 10 |
| Tetanus..... | 1 |
| Tuberculosis..... | 8 |
| Typhoid fever..... | 7 |
| Typhus fever..... | 1 |
| Whooping cough..... | 8 |

GEORGIA

| | |
|-------------------------|----|
| Dengue..... | 1 |
| Diphtheria..... | 12 |
| Dysentery..... | 9 |
| Hookworm disease..... | 4 |
| Influenza..... | 15 |
| Malaria..... | 46 |
| Measles..... | 4 |
| Mumps..... | 2 |
| Paratyphoid fever..... | 5 |
| Pellagra..... | 1 |
| Pneumonia..... | 5 |
| Polioimyelitis..... | 1 |
| Rabies..... | 1 |
| Scarlet fever..... | 3 |
| Septic sore throat..... | 30 |
| Smallpox..... | 5 |
| Tuberculosis..... | 4 |
| Typhoid fever..... | 68 |
| Whooping cough..... | 22 |

IDAHO

| | |
|---------------------|----|
| Chicken pox..... | 1 |
| Measles..... | 1 |
| Mumps..... | 1 |
| Scarlet fever..... | 5 |
| Smallpox..... | 2 |
| Typhoid fever..... | 1 |
| Whooping cough..... | 10 |

ILLINOIS

| | |
|---------------------------|---|
| Cerebrospinal meningitis: | |
| Cook County..... | 2 |
| Moultrie County..... | 1 |
| Peoria County..... | 1 |

ILLINOIS—continued

| | Cases |
|-------------------------|-------|
| Chicken pox..... | 17 |
| Diphtheria..... | 60 |
| Influenza..... | 37 |
| Lethargic encephalitis: | |
| Cook County..... | 1 |
| Henry County..... | 1 |
| Jackson County..... | 1 |
| Lawrence County..... | 1 |
| Lee County..... | 1 |
| Perry County..... | 1 |
| Schuyler County..... | 1 |
| Stephenson County..... | 1 |
| Measles..... | 50 |
| Mumps..... | 23 |
| Pneumonia..... | 156 |
| Polioimyelitis: | |
| Iroquois County..... | 1 |
| McLean County..... | 1 |
| Peoria County..... | 1 |
| Tazewell County..... | 2 |
| Scarlet fever..... | 59 |
| Smallpox..... | 3 |
| Tuberculosis..... | 425 |
| Typhoid fever..... | 41 |
| Whooping cough..... | 185 |

INDIANA

| | |
|-------------------------------|----|
| Cerebrospinal meningitis..... | 1 |
| Chicken pox..... | 1 |
| Diphtheria..... | 15 |
| Influenza..... | 9 |
| Measles..... | 13 |
| Scarlet fever..... | 15 |
| Smallpox..... | 3 |
| Tuberculosis..... | 50 |
| Typhoid fever..... | 21 |
| Whooping cough..... | 15 |

IOWA

| | |
|---------------------|----|
| Chicken pox..... | 3 |
| Diphtheria..... | 12 |
| Measles..... | 1 |
| Mumps..... | 2 |
| Pneumonia..... | 1 |
| Scarlet fever..... | 8 |
| Smallpox..... | 2 |
| Tuberculosis..... | 4 |
| Typhoid fever..... | 6 |
| Whooping cough..... | 6 |

KANSAS

| | |
|---------------------------------------|----|
| Cerebrospinal meningitis—Mayetta..... | 1 |
| Chicken pox..... | 2 |
| Diphtheria..... | 10 |
| Influenza..... | 4 |
| Malaria..... | 7 |
| Measles..... | 5 |
| Pneumonia..... | 17 |
| Polioimyelitis: | |
| Hutchinson..... | 3 |
| Jennings..... | 1 |
| Norcatur..... | 1 |
| Spearville..... | 1 |
| Stafford..... | 1 |
| Scarlet fever..... | 29 |
| Smallpox..... | 2 |

KANSAS—continued

| | Cases |
|---------------------|-------|
| Tetanus..... | 1 |
| Tuberculosis..... | 50 |
| Typhoid fever..... | 22 |
| Whooping cough..... | 38 |

LOUISIANA

| | |
|-----------------------------|----|
| Diphtheria..... | 14 |
| Influenza..... | 7 |
| Lethargic encephalitis..... | 1 |
| Malaria..... | 16 |
| Pneumonia..... | 20 |
| Polioymyelitis..... | 1 |
| Scarlet fever..... | 7 |
| Smallpox..... | 5 |
| Tuberculosis..... | 35 |
| Typhoid fever..... | 29 |

MAINE

| | |
|-----------------------|----|
| Chicken pox..... | 9 |
| Diphtheria..... | 1 |
| Influenza..... | 6 |
| Measles..... | 26 |
| Pneumonia..... | 2 |
| Scarlet fever..... | 20 |
| Tuberculosis..... | 5 |
| Typhoid fever..... | 14 |
| Vincent's angina..... | 2 |
| Whooping cough..... | 11 |

MARYLAND¹

| | |
|-------------------------------|----|
| Cerebrospinal meningitis..... | 3 |
| Chicken pox..... | 4 |
| Diphtheria..... | 18 |
| Dysentery..... | 16 |
| Influenza..... | 3 |
| Lethargic encephalitis..... | 5 |
| Malaria..... | 1 |
| Measles..... | 7 |
| Mumps..... | 3 |
| Paratyphoid fever..... | 3 |
| Pneumonia (broncho)..... | 7 |
| Pneumonia (lobar)..... | 11 |
| Polioymyelitis..... | 5 |
| Scarlet fever..... | 9 |
| Tetanus..... | 1 |
| Trachoma..... | 1 |
| Tuberculosis..... | 42 |
| Typhoid fever..... | 72 |
| Whooping cough..... | 72 |

MASSACHUSETTS

| | |
|-----------------------------------|----|
| Cerebrospinal meningitis..... | 1 |
| Chicken pox..... | 10 |
| Conjunctivitis (suppurative)..... | 6 |
| Diphtheria..... | 46 |
| Dysentery..... | 1 |
| German measles..... | 4 |
| Lethargic encephalitis..... | 1 |
| Measles..... | 17 |
| Mumps..... | 24 |
| Ophthalmia neonatorum..... | 34 |
| Pellagra..... | 1 |
| Pneumonia (lobar)..... | 16 |
| Polioymyelitis..... | 10 |
| Scarlet fever..... | 54 |
| Septic sore throat..... | 2 |

MASSACHUSETTS—continued

| | Cases |
|---------------------------------|-------|
| Trachoma..... | 1 |
| Trichinosis..... | 1 |
| Tuberculosis (pulmonary)..... | 104 |
| Tuberculosis (other forms)..... | 35 |
| Typhoid fever..... | 14 |
| Whooping cough..... | 81 |

MICHIGAN

| | |
|---------------------|-----|
| Diphtheria..... | 85 |
| Measles..... | 26 |
| Pneumonia..... | 68 |
| Scarlet fever..... | 72 |
| Smallpox..... | 1 |
| Tuberculosis..... | 240 |
| Typhoid fever..... | 18 |
| Whooping cough..... | 174 |

MINNESOTA

| | |
|-----------------------------|----|
| Chicken pox..... | 7 |
| Diphtheria..... | 28 |
| Influenza..... | 1 |
| Lethargic encephalitis..... | 2 |
| Measles..... | 20 |
| Polioymyelitis..... | 3 |
| Scarlet fever..... | 59 |
| Smallpox..... | 3 |
| Trachoma..... | 1 |
| Tuberculosis..... | 33 |
| Typhoid fever..... | 7 |
| Whooping cough..... | 27 |

MISSISSIPPI

| | |
|---------------------|----|
| Diphtheria..... | 22 |
| Polioymyelitis..... | 1 |
| Scarlet fever..... | 6 |
| Smallpox..... | 2 |
| Typhoid fever..... | 51 |

MISSOURI

(Exclusive of Kansas City)

| | |
|---------------------|----|
| Chicken pox..... | 2 |
| Diphtheria..... | 15 |
| Malaria..... | 1 |
| Measles..... | 7 |
| Mumps..... | 11 |
| Polioymyelitis..... | 1 |
| Scarlet fever..... | 37 |
| Smallpox..... | 1 |
| Trachoma..... | 5 |
| Tuberculosis..... | 32 |
| Typhoid fever..... | 39 |
| Whooping cough..... | 27 |

MONTANA

| | |
|-----------------------------------|----|
| Chicken pox..... | 1 |
| Diphtheria..... | 2 |
| Measles..... | 5 |
| Mumps..... | 1 |
| Rocky Mountain spotted fever..... | 3 |
| Scarlet fever..... | 13 |
| Smallpox..... | 3 |
| Tetanus..... | 1 |
| Tuberculosis..... | 2 |
| Typhoid fever..... | 4 |

¹ Week ended Friday.

| NEBRASKA | Cases |
|-------------------------|-------|
| Diphtheria..... | 1 |
| Measles..... | 3 |
| Poliomyelitis..... | 1 |
| Scarlet fever..... | 2 |
| Septic sore throat..... | 2 |
| Smallpox..... | 2 |
| Tuberculosis..... | 1 |
| Typhoid fever..... | 4 |
| Whooping cough..... | 41 |

| NEW JERSEY | Cases |
|------------------------|-------|
| Chicken pox..... | 8 |
| Diphtheria..... | 33 |
| Dysentery..... | 1 |
| Influenza..... | 7 |
| Measles..... | 15 |
| Paratyphoid fever..... | 1 |
| Pneumonia..... | 18 |
| Poliomyelitis..... | 1 |
| Scarlet fever..... | 15 |
| Typhoid fever..... | 24 |
| Whooping cough..... | 134 |

| NEW MEXICO | Cases |
|--------------------------|-------|
| Chicken pox..... | 2 |
| Diphtheria..... | 1 |
| Measles..... | 2 |
| Mumps..... | 4 |
| Pneumonia..... | 1 |
| Rabies (in animals)..... | 3 |
| Tuberculosis..... | 21 |
| Typhoid fever..... | 6 |
| Whooping cough..... | 6 |

| NEW YORK | Cases |
|-------------------------------|-------|
| (Exclusive of New York City) | |
| Cerebrospinal meningitis..... | 1 |
| Chicken pox..... | 30 |
| Diphtheria..... | 34 |
| German measles..... | 8 |
| Malaria..... | 12 |
| Measles..... | 76 |
| Mumps..... | 18 |
| Ophthalmia neonatorum..... | 1 |
| Paratyphoid fever..... | 1 |
| Pneumonia..... | 38 |
| Poliomyelitis..... | 52 |
| Scarlet fever..... | 37 |
| Tetanus..... | 1 |
| Typhoid fever..... | 22 |
| Vincent's angina..... | 6 |
| Whooping cough..... | 104 |

| NORTH CAROLINA | Cases |
|----------------------------|-------|
| Chicken pox..... | 4 |
| Diphtheria..... | 56 |
| Dysentery (bacillary)..... | 7 |
| German measles..... | 2 |
| Malaria..... | 23 |
| Measles..... | 24 |
| Ophthalmia neonatorum..... | 1 |
| Poliomyelitis..... | 12 |
| Scarlet fever..... | 35 |
| Septic sore throat..... | 4 |
| Smallpox..... | 17 |
| Typhoid fever..... | 107 |
| Whooping cough..... | 221 |

¹ Deaths.

| OKLAHOMA | Cases |
|--|-------|
| (Exclusive of Oklahoma City and Tulsa) | |
| Diphtheria..... | 10 |
| Influenza..... | 18 |
| Malaria..... | 120 |
| Measles..... | 6 |
| Pellagra..... | 12 |
| Pneumonia..... | 7 |
| Poliomyelitis—Hughes County..... | 1 |
| Scarlet fever..... | 4 |
| Typhoid fever..... | 94 |
| Whooping cough..... | 14 |

| OREGON | Cases |
|-------------------------|-------|
| Chicken pox..... | 3 |
| Diphtheria..... | 10 |
| Dysentery..... | 1 |
| Influenza..... | 14 |
| Measles..... | 6 |
| Mumps..... | 3 |
| Pneumonia..... | 15 |
| Poliomyelitis..... | 1 |
| Scarlet fever..... | 8 |
| Septic sore throat..... | 1 |
| Smallpox..... | 9 |
| Tuberculosis..... | 10 |
| Typhoid fever..... | 6 |
| Whooping cough..... | 12 |

| PENNSYLVANIA | Cases |
|---|-------|
| Chicken pox..... | 53 |
| Diphtheria..... | 117 |
| German measles..... | 12 |
| Impetigo contagiosa..... | 20 |
| Measles..... | 123 |
| Mumps..... | 8 |
| Ophthalmia neonatorum—Philadelphia..... | 8 |
| Pneumonia..... | 10 |
| Poliomyelitis: | |
| Brothers Valley Township ¹ | 1 |
| Chambersburg..... | 1 |
| Johnstown..... | 2 |
| Philadelphia..... | 2 |
| Scarlet fever..... | 127 |
| Tetanus: | |
| Franklin..... | 1 |
| North Whitehall Township ¹ | 1 |
| St. Clair..... | 1 |
| Trachoma—Philadelphia..... | 1 |
| Tuberculosis..... | 100 |
| Typhoid fever..... | 84 |
| Whooping cough..... | 389 |

| RHODE ISLAND | Cases |
|----------------------------|-------|
| Diphtheria..... | 1 |
| German measles..... | 1 |
| Influenza..... | 4 |
| Measles..... | 1 |
| Pneumonia..... | 1 |
| Poliomyelitis—Newport..... | 1 |
| Scarlet fever..... | 3 |
| Tuberculosis..... | 10 |
| Typhoid fever..... | 1 |
| Whooping cough..... | 13 |

¹ County not specified

| SOUTH DAKOTA | | WASHINGTON | |
|-----------------------------------|-------|--|-------|
| | Cases | | Cases |
| Diphtheria..... | 1 | Cerebrospinal meningitis..... | 1 |
| Measles..... | 2 | Chicken pox..... | 10 |
| Pneumonia..... | 1 | Diphtheria..... | 8 |
| Scarlet fever..... | 11 | German measles..... | 4 |
| Smallpox..... | 3 | Measles..... | 6 |
| Tuberculosis..... | 2 | Mumps..... | 1 |
| Typhoid fever..... | 5 | Pneumonia..... | 1 |
| Whooping cough..... | 4 | Scarlet fever..... | 14 |
| | | Smallpox..... | 6 |
| | | Tuberculosis..... | 17 |
| | | Typhoid fever..... | 25 |
| | | Whooping cough..... | 19 |
| TENNESSEE | | WEST VIRGINIA | |
| Chicken pox..... | 4 | Chicken pox..... | 1 |
| Diphtheria..... | 17 | Diphtheria..... | 16 |
| Dysentery..... | 2 | Influenza..... | 20 |
| Influenza..... | 4 | Measles..... | 22 |
| Malaria..... | 47 | Scarlet fever..... | 20 |
| Measles..... | 6 | Smallpox..... | 3 |
| Pellagra..... | 5 | Tuberculosis..... | 8 |
| Pneumonia..... | 3 | Typhoid fever..... | 24 |
| Poliomyelitis: | | Whooping cough..... | 30 |
| Davidson County..... | 1 | | |
| Franklin County..... | 1 | | |
| Overton County..... | 1 | | |
| Scarlet fever..... | 17 | | |
| Smallpox..... | 1 | | |
| Tuberculosis..... | 28 | | |
| Typhoid fever..... | 206 | | |
| Whooping cough..... | 13 | | |
| TEXAS | | WISCONSIN | |
| Chicken pox..... | 5 | Milwaukee: | |
| Diphtheria..... | 13 | Chicken pox..... | 1 |
| Influenza..... | 19 | Diphtheria..... | 5 |
| Measles..... | 3 | German measles..... | 1 |
| Mumps..... | 1 | Measles..... | 8 |
| Paratyphoid fever..... | 2 | Mumps..... | 2 |
| Pellagra..... | 1 | Pneumonia..... | 2 |
| Pneumonia..... | 2 | Poliomyelitis..... | 1 |
| Poliomyelitis..... | 1 | Scarlet fever..... | 8 |
| Scarlet fever..... | 11 | Tuberculosis..... | 8 |
| Smallpox..... | 2 | Typhoid fever..... | 1 |
| Tuberculosis..... | 33 | Whooping cough..... | 66 |
| Typhoid fever..... | 20 | Scattering: | |
| Whooping cough..... | 30 | Chicken pox..... | 13 |
| | | Diphtheria..... | 13 |
| | | German measles..... | 4 |
| | | Influenza..... | 2 |
| | | Measles..... | 91 |
| | | Mumps..... | 6 |
| | | Pneumonia..... | 3 |
| | | Poliomyelitis..... | 1 |
| | | Scarlet fever..... | 38 |
| | | Smallpox..... | 6 |
| | | Tuberculosis..... | 37 |
| | | Typhoid fever..... | 11 |
| | | Whooping cough..... | 123 |
| UTAH | | WYOMING | |
| Chicken pox..... | 3 | Cerebrospinal meningitis—Natrona County... | 1 |
| Diphtheria..... | 13 | Chicken pox..... | 1 |
| Measles..... | 3 | Scarlet fever..... | 2 |
| Pneumonia..... | 4 | Smallpox..... | 1 |
| Poliomyelitis—Salt Lake City..... | 1 | Typhoid fever..... | 1 |
| Scarlet fever..... | 2 | Whooping cough..... | 2 |
| Typhoid fever..... | 1 | | |
| Whooping cough..... | 27 | | |
| VERMONT | | | |
| Chicken pox..... | 3 | | |
| Diphtheria..... | 3 | | |
| Measles..... | 3 | | |
| Mumps..... | 1 | | |
| Whooping cough..... | 19 | | |

Reports for Week Ended August 28, 1926

| CALIFORNIA | | Cases | DISTRICT OF COLUMBIA—continued | | Cases |
|---|-----|-------|--------------------------------|--|-------|
| Cerebrospinal meningitis—San Bernardino County..... | 1 | | Typhoid fever..... | | 3 |
| Chicken pox..... | 33 | | Whooping cough..... | | 13 |
| Diphtheria..... | 75 | | NORTH DAKOTA | | |
| Influenza..... | 6 | | Chicken pox..... | | 2 |
| Lethargic encephalitis: | | | Diphtheria..... | | 1 |
| Los Angeles County..... | 1 | | German measles..... | | 3 |
| Santa Ana..... | 1 | | Pneumonia..... | | 3 |
| Measles..... | 141 | | Scarlet fever..... | | 8 |
| Mumps..... | 73 | | Tuberculosis..... | | 3 |
| Poliomyelitis: | | | Typhoid fever..... | | 2 |
| Los Angeles..... | 1 | | Whooping cough..... | | 20 |
| Los Angeles County..... | 1 | | SOUTH CAROLINA | | |
| California ¹ | 1 | | Chicken pox..... | | 3 |
| Scarlet fever..... | 65 | | Dengue..... | | 32 |
| Smallpox..... | 4 | | Diphtheria..... | | 28 |
| Tuberculosis..... | 171 | | Hookworm disease..... | | 28 |
| Typhoid fever..... | 23 | | Influenza..... | | 78 |
| Whooping cough..... | 54 | | Malaria..... | | 382 |
| DISTRICT OF COLUMBIA | | | Paratyphoid fever..... | | 21 |
| Chicken pox..... | 1 | | Pellagra..... | | 64 |
| Diphtheria..... | 7 | | Poliomyelitis..... | | 2 |
| Influenza..... | 1 | | Scarlet fever..... | | 9 |
| Pneumonia..... | 2 | | Smallpox..... | | 8 |
| Poliomyelitis..... | 1 | | Tuberculosis..... | | 48 |
| Scarlet fever..... | 14 | | Typhoid fever..... | | 118 |
| Tuberculosis..... | 14 | | Whooping cough..... | | 34 |

¹ Place not specified.

SUMMARY OF MONTHLY REPORTS FROM STATES

The following summary of monthly State reports is published weekly and covers only those States from which reports are received during the current week:

| State | Cerebrospinal meningitis | Diphtheria | Influenza | Malaria | Measles | Pellagra | Poliomyelitis | Scarlet fever | Smallpox | Typhoid fever |
|-------------------|--------------------------|------------|-----------|---------|---------|----------|---------------|---------------|----------|---------------|
| <i>July, 1926</i> | | | | | | | | | | |
| Florida..... | 3 | 66 | 301 | 33 | 55 | 7 | 3 | 14 | 48 | 64 |
| Georgia..... | 2 | 26 | 66 | 216 | 124 | 56 | 1 | 13 | 29 | 414 |
| Iowa..... | 2 | 37 | | | 72 | | 0 | 76 | 59 | 10 |
| Louisiana..... | 0 | 25 | 166 | 126 | 1 | 37 | 2 | 24 | 5 | 188 |
| Maryland..... | | 45 | 9 | 4 | 360 | 1 | 11 | 97 | 0 | 68 |
| Virginia..... | 5 | 64 | 280 | 175 | 638 | 21 | 22 | 68 | 44 | 188 |
| Washington..... | 12 | 128 | 8 | | 183 | | 3 | 118 | 85 | 36 |

GENERAL CURRENT SUMMARY AND WEEKLY REPORTS FROM CITIES

Diphtheria.—For the week ended August 21, 1926, 37 States reported 655 cases of diphtheria. For the week ended August 22, 1925, the same States reported 879 cases of this disease. Ninety-eight cities, situated in all parts of the country and having an aggregate population of more than 30,200,000, reported 397 cases of diphtheria for the week ended August 21, 1926. Last year for the corresponding week they reported 388 cases. The estimated expectancy for these cities was 541 cases. The estimated expectancy is based on the experience of the last nine years, excluding epidemics.

Measles.—Thirty-five States reported 975 cases of measles for the week ended August 21, 1926, and 401 cases of this disease for the week ended August 22, 1925. Ninety-eight cities reported 238 cases of measles for the week this year, and 170 cases last year.

Poliomyelitis.—The health officers of 37 States reported 99 cases of poliomyelitis for the week ended August 21, 1926. The same States reported 296 cases for the week ended August 22, 1925.

Scarlet fever.—Scarlet fever was reported for the week as follows: Thirty-seven States—this year, 858 cases; last year, 705 cases; 98 cities—this year, 281 cases; last year, 285 cases; estimated expectancy, 238 cases.

Smallpox.—For the week ended August 21, 1926, 37 States reported 129 cases of smallpox. Last year for the corresponding week they reported 135 cases. Ninety-eight cities reported smallpox for the week as follows: 1926, 12 cases; 1925, 29 cases; estimated expectancy, 23 cases. No deaths from smallpox were reported by these cities for the week this year.

Typhoid fever.—One thousand two hundred and fifty-one cases of typhoid fever were reported for the week ended August 21, 1926, by 36 States. For the corresponding week of 1925, the same States reported 1,319 cases of this disease. Ninety-eight cities reported 237 cases of typhoid fever for the week this year and 314 cases for the corresponding week last year. The estimated expectancy for these cities was 235 cases.

Influenza and pneumonia.—Deaths from influenza and pneumonia were reported for the week by 94 cities, with a population of nearly 29,700,000, as follows: 1926, 324 deaths; 1925, 309 deaths.

City reports for week ended August 21, 1926

The "estimated expectancy" given for diphtheria, poliomyelitis, scarlet fever, smallpox, and typhoid fever is the result of an attempt to ascertain from previous occurrence how many cases of the disease under consideration may be expected to occur during a certain week in the absence of epidemics. It is based on reports to the Public Health Service during the past nine years. It is in most instances the median number of cases reported in the corresponding week of the preceding years. When the reports include several epidemics or when for other reasons the median is unsatisfactory, the epidemic periods are excluded and the estimated expectancy is the mean number of cases reported for the week during nonepidemic years.

If reports have not been received for the full nine years, data are used for as many years as possible, but no year earlier than 1917 is included. In obtaining the estimated expectancy the figures are smoothed when necessary to avoid abrupt deviations from the usual trend. For some of the diseases given in the table the available data were not sufficient to make it practicable to compute the estimated expectancy.

| Division, State, and city | Population July 1, 1925, estimated | Chicken pox, cases reported | Diphtheria | | Influenza | | Measles, cases reported | Mumps, cases reported | Pneumonia, deaths reported |
|---------------------------|------------------------------------|-----------------------------|-----------------------------|----------------|----------------|-----------------|-------------------------|-----------------------|----------------------------|
| | | | Cases, estimated expectancy | Cases reported | Cases reported | Deaths reported | | | |
| NEW ENGLAND | | | | | | | | | |
| Maine: | | | | | | | | | |
| Portland..... | 75,333 | 0 | 1 | 0 | 0 | 0 | 1 | 1 | 4 |
| New Hampshire: | | | | | | | | | |
| Concord..... | 22,546 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Manchester..... | 83,097 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 1 |
| Vermont: | | | | | | | | | |
| Barre..... | 10,608 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Massachusetts: | | | | | | | | | |
| Boston..... | 779,620 | 11 | 31 | 6 | 1 | 0 | 15 | 14 | 9 |
| Fall River..... | 128,993 | 2 | 2 | 3 | 0 | 0 | 0 | 0 | 0 |
| Springfield..... | 142,665 | 0 | 2 | 0 | 0 | 0 | 0 | 0 | 0 |
| Worcester..... | 190,757 | 0 | 2 | 4 | 0 | 0 | 0 | 0 | 0 |
| Rhode Island: | | | | | | | | | |
| Pawtucket..... | 69,760 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Providence..... | 267,918 | 0 | 3 | 2 | 0 | 0 | 0 | 0 | 1 |
| Connecticut: | | | | | | | | | |
| Bridgeport..... | (1) | 0 | 4 | 3 | 0 | 0 | 0 | 0 | 0 |
| Hartford..... | 160,197 | 1 | 3 | 1 | 0 | 0 | 2 | 0 | 3 |
| New Haven..... | 178,927 | 0 | 2 | 1 | 0 | 0 | 4 | 0 | 0 |
| MIDDLE ATLANTIC | | | | | | | | | |
| New York: | | | | | | | | | |
| Buffalo..... | 538,010 | 0 | 11 | 10 | 0 | 1 | 3 | 5 | |
| New York..... | 5,873,356 | 35 | 107 | 69 | 6 | 14 | 14 | 80 | |
| Rochester..... | 316,786 | 2 | 4 | 2 | 0 | 1 | 2 | 2 | |
| Syracuse..... | 182,003 | 0 | 3 | 1 | 0 | 10 | 0 | 0 | |
| New Jersey: | | | | | | | | | |
| Camden..... | 128,642 | 0 | 2 | 7 | 0 | 2 | 0 | 1 | |
| Newark..... | 452,513 | 4 | 7 | 2 | 2 | 8 | 2 | 4 | |
| Trenton..... | 132,020 | 0 | 2 | 1 | 0 | 0 | 0 | 0 | |
| Pennsylvania: | | | | | | | | | |
| Philadelphia..... | 1,979,364 | 12 | 33 | 23 | 0 | 7 | 0 | 15 | |
| Pittsburgh..... | 631,563 | 2 | 15 | 4 | 2 | 11 | 0 | 10 | |
| Reading..... | 112,707 | 3 | 2 | 0 | 0 | 1 | 0 | 0 | |
| EAST NORTH CENTRAL | | | | | | | | | |
| Ohio: | | | | | | | | | |
| Cincinnati..... | 409,333 | 0 | 6 | 3 | 0 | 3 | 1 | 6 | |
| Cleveland..... | 936,485 | 10 | 19 | 35 | 0 | 5 | 1 | 6 | |
| Columbus..... | 270,836 | 0 | 2 | 4 | 0 | 2 | 0 | 1 | |
| Toledo..... | 287,380 | 1 | 5 | 1 | 0 | 2 | 0 | 4 | |
| Indiana: | | | | | | | | | |
| Fort Wayne..... | 97,846 | 0 | 1 | 0 | 0 | 0 | 0 | 2 | |
| Indianapolis..... | 355,819 | 0 | 5 | 1 | 0 | 0 | 0 | 3 | |
| South Bend..... | 80,091 | 0 | 0 | 2 | 0 | 3 | 0 | 1 | |
| Terre Haute..... | 71,071 | 0 | 1 | 0 | 1 | 0 | 0 | 0 | |
| Illinois: | | | | | | | | | |
| Chicago..... | 2,995,239 | 22 | 59 | 28 | 2 | 42 | 5 | 15 | |
| Peoria..... | 81,564 | 0 | 1 | 1 | 0 | 1 | 0 | 1 | |
| Springfield..... | 63,923 | 1 | 1 | 0 | 0 | 0 | 0 | 0 | |
| Michigan: | | | | | | | | | |
| Detroit..... | 1,245,824 | 8 | 25 | 45 | 0 | 10 | 2 | 5 | |
| Flint..... | 130,316 | 0 | 4 | 1 | 0 | 0 | 0 | 1 | |
| Grand Rapids..... | 153,698 | 0 | 2 | 0 | 0 | 2 | 0 | 1 | |

¹ No estimate made.

City reports for week ended August 21, 1926—Continued

| Division, State, and city | Population July 1, 1925, estimated | Chick- en pox, cases re- ported | Diphtheria | | Influenza | | Meas- les, cases re- ported | Mumps, cases re- ported | Pneu- monia, deaths re- ported |
|----------------------------------|------------------------------------|--|---------------------------------------|---------------------|---------------------|----------------------|--------------------------------------|-------------------------------|---|
| | | | Cases, esti- mated expec- tancy | Cases re- ported | Cases re- ported | Deaths re- ported | | | |
| EAST NORTH CENTRAL— continued | | | | | | | | | |
| Wisconsin: | | | | | | | | | |
| Kenosha..... | 50,891 | 0 | 1 | 0 | 0 | 0 | 17 | 0 | 1 |
| Madison..... | 46,385 | | 1 | | | | | | |
| Milwaukee..... | 509,192 | 4 | 10 | 4 | 0 | 0 | 17 | 5 | 3 |
| Racine..... | 67,707 | 2 | 1 | 0 | 0 | 0 | 4 | 0 | 1 |
| Superior..... | 39,671 | 0 | 0 | 4 | 0 | 0 | 0 | 0 | 0 |
| WEST NORTH CENTRAL | | | | | | | | | |
| Minnesota: | | | | | | | | | |
| Duluth..... | 110,502 | 0 | 2 | 0 | 0 | 0 | 3 | 0 | 3 |
| Minneapolis..... | 425,455 | 3 | 12 | 15 | 0 | 0 | 0 | 0 | 4 |
| St. Paul..... | 246,001 | 0 | 11 | 6 | 0 | 1 | 1 | 0 | 8 |
| Iowa: | | | | | | | | | |
| Davenport..... | 52,469 | 0 | 1 | 0 | 0 | | 0 | 0 | |
| Sioux City..... | 76,411 | 0 | 1 | 0 | 0 | | 0 | 0 | |
| Waterloo..... | 36,771 | 0 | 0 | 0 | 0 | | 2 | 0 | |
| Missouri: | | | | | | | | | |
| Kansas City..... | 367,481 | 0 | 3 | 0 | 0 | 0 | 4 | 0 | 4 |
| St. Joseph..... | 78,342 | 0 | 0 | 0 | 0 | 0 | 1 | 0 | 2 |
| St. Louis..... | 821,543 | 3 | 18 | 20 | 0 | 0 | 0 | 0 | |
| North Dakota: | | | | | | | | | |
| Fargo..... | 26,403 | 0 | 1 | 0 | 0 | 0 | 0 | 0 | 0 |
| Grand Forks..... | 14,811 | 0 | 0 | 0 | 0 | | 0 | 0 | |
| South Dakota: | | | | | | | | | |
| Aberdeen..... | 15,036 | 0 | 0 | 0 | 0 | | 0 | 0 | |
| Sioux Falls..... | 30,127 | | 0 | | | | | | |
| Nebraska: | | | | | | | | | |
| Lincoln..... | 60,941 | 0 | 1 | 0 | 0 | 0 | 0 | 1 | 0 |
| Omaha..... | 211,768 | 0 | 5 | 0 | 0 | 0 | 1 | 0 | 1 |
| Kansas: | | | | | | | | | |
| Topeka..... | 55,411 | 0 | 1 | 0 | 0 | 0 | 1 | 0 | 1 |
| Wichita..... | 88,367 | 0 | 1 | 0 | 0 | 0 | 1 | 0 | 0 |
| SOUTH ATLANTIC | | | | | | | | | |
| Delaware: | | | | | | | | | |
| Wilmington..... | 122,049 | 0 | 1 | 0 | 0 | 0 | 0 | 1 | 0 |
| Maryland: | | | | | | | | | |
| Baltimore..... | 766,296 | 4 | 12 | 8 | 2 | 0 | 0 | 3 | 18 |
| Cumberland..... | 33,741 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Frederick..... | 12,035 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| District of Columbia: | | | | | | | | | |
| Washington..... | 497,906 | 0 | 3 | 7 | 0 | 0 | 3 | 0 | 8 |
| Virginia: | | | | | | | | | |
| Lynchburg..... | 30,395 | 0 | 1 | 1 | 0 | 0 | 0 | 0 | 1 |
| Norfolk..... | (1) | 0 | 1 | 0 | 0 | 0 | 0 | 0 | 2 |
| Richmond..... | 186,463 | 0 | 5 | 12 | 0 | 0 | 7 | 0 | 2 |
| Roanoke..... | 58,208 | 0 | 2 | 0 | 0 | 0 | 0 | 0 | 0 |
| West Virginia: | | | | | | | | | |
| Charleston..... | 49,019 | 0 | 1 | 0 | 0 | 1 | 0 | 0 | 0 |
| Huntington..... | 63,485 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 1 |
| Wheeling..... | 56,208 | 0 | 0 | 0 | 0 | 0 | 1 | 0 | 1 |
| North Carolina: | | | | | | | | | |
| Raleigh..... | 30,371 | 0 | 0 | 0 | 0 | 0 | 1 | 0 | 2 |
| Wilmington..... | 37,661 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 1 |
| Winston-Salem..... | 69,031 | 0 | 1 | 0 | 0 | 0 | 1 | 0 | 3 |
| South Carolina: | | | | | | | | | |
| Charleston..... | 73,125 | 0 | 1 | 1 | 0 | 0 | 0 | 0 | 0 |
| Columbia..... | 41,225 | 0 | 1 | 1 | 0 | 0 | 0 | 0 | 0 |
| Greenville..... | 27,311 | 0 | 1 | 0 | 0 | 0 | 0 | 0 | 0 |
| Georgia: | | | | | | | | | |
| Atlanta..... | (1) | 1 | 2 | 1 | 2 | 0 | 0 | 0 | 3 |
| Brunswick..... | 16,800 | 0 | 0 | 0 | 2 | 0 | 0 | 0 | 2 |
| Savannah..... | 93,134 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Florida: | | | | | | | | | |
| Miami..... | 69,754 | 0 | | 1 | 0 | 0 | 1 | 0 | 2 |
| St. Petersburg..... | 26,847 | | 0 | | | 0 | | | 0 |
| Tampa..... | 94,743 | 0 | 1 | 1 | 0 | 0 | 6 | 0 | 3 |

1 No estimate made.

City reports for week ended August 21, 1926—Continued

| Division, State, and city | Population July 1, 1925, estimated | Chick- en pox, cases re- ported | Diphtheria | | Influenza | | Meas- les, cases re- ported | Mumps, cases re- ported | Pneu- monia, deaths re- ported |
|---------------------------|------------------------------------|--|---------------------------------------|---------------------|---------------------|----------------------|--------------------------------------|-------------------------------|---|
| | | | Cases, esti- mated expec- tancy | Cases re- ported | Cases re- ported | Deaths re- ported | | | |
| EAST SOUTH CENTRAL | | | | | | | | | |
| Kentucky: | | | | | | | | | |
| Covington..... | 58,309 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Louisville..... | 305,935 | 0 | 3 | 0 | 0 | 0 | 1 | 0 | 3 |
| Tennessee: | | | | | | | | | |
| Memphis..... | 174,533 | 0 | 3 | 0 | 0 | 0 | 0 | 0 | 0 |
| Nashville..... | 136,220 | 0 | 1 | 3 | 0 | 0 | 0 | 0 | 2 |
| Alabama: | | | | | | | | | |
| Birmingham..... | 205,670 | 1 | 2 | 1 | 5 | 0 | 6 | 1 | 1 |
| Mobile..... | 65,955 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 1 |
| Montgomery..... | 46,481 | 0 | 1 | 0 | 0 | 0 | 0 | 0 | 0 |
| WEST SOUTH CENTRAL | | | | | | | | | |
| Arkansas: | | | | | | | | | |
| Fort Smith..... | 31,643 | | 1 | | | | | | |
| Little Rock..... | 74,216 | 0 | 1 | 1 | 0 | 0 | 0 | 0 | 2 |
| Louisiana: | | | | | | | | | |
| New Orleans..... | 414,493 | 0 | 6 | 11 | 4 | 5 | 1 | 0 | 2 |
| Shreveport..... | 57,857 | 0 | 0 | 1 | 0 | 0 | 0 | 0 | 1 |
| Oklahoma: | | | | | | | | | |
| Oklahoma City..... | (1) | 0 | 1 | 0 | 0 | 0 | 0 | 0 | 3 |
| Texas: | | | | | | | | | |
| Dallas..... | 194,450 | 0 | 3 | 1 | 0 | 0 | 1 | 1 | 4 |
| Galveston..... | 48,375 | 0 | 1 | 0 | 0 | 0 | 0 | 0 | 0 |
| Houston..... | 164,934 | 0 | 1 | 0 | 0 | 0 | 0 | 0 | 1 |
| San Antonio..... | 198,069 | 0 | 1 | 1 | 0 | 1 | 0 | 0 | 5 |
| MOUNTAIN | | | | | | | | | |
| Montana: | | | | | | | | | |
| Billings..... | 17,971 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 1 |
| Great Falls..... | 20,883 | 0 | 1 | 0 | 0 | 0 | 0 | 0 | 0 |
| Helena..... | 12,037 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Missoula..... | 12,668 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Idaho: | | | | | | | | | |
| Boise..... | 23,042 | 0 | 0 | 1 | 0 | 0 | 0 | 0 | 0 |
| Colorado: | | | | | | | | | |
| Denver..... | 280,911 | 3 | 8 | 11 | | 0 | 1 | 1 | 4 |
| Pueblo..... | 43,787 | 0 | 3 | 1 | 0 | 0 | 0 | 0 | 0 |
| New Mexico: | | | | | | | | | |
| Albuquerque..... | 21,000 | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 0 |
| Arizona: | | | | | | | | | |
| Phoenix..... | 38,669 | 0 | 0 | 1 | 0 | 0 | 0 | 0 | 1 |
| Utah: | | | | | | | | | |
| Salt Lake City..... | 130,948 | 1 | 2 | 3 | 0 | 0 | 1 | 0 | 4 |
| Nevada: | | | | | | | | | |
| Reno..... | 12,665 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| PACIFIC | | | | | | | | | |
| Washington: | | | | | | | | | |
| Seattle..... | (1) | 11 | 3 | 1 | 0 | | 4 | 4 | |
| Spokane..... | 108,897 | 1 | 2 | 2 | 0 | | 10 | 0 | |
| Tacoma..... | 104,455 | 0 | 1 | 2 | 0 | 1 | 0 | 0 | 1 |
| Oregon: | | | | | | | | | |
| Portland..... | 282,383 | 0 | 4 | 6 | 0 | 0 | 8 | 2 | 4 |
| California: | | | | | | | | | |
| Los Angeles..... | (1) | 11 | 24 | 12 | 3 | 0 | 1 | 2 | 12 |
| Sacramento..... | 72,260 | 1 | 2 | 0 | 0 | 0 | 1 | 1 | 0 |
| San Francisco..... | 557,530 | 6 | 13 | 6 | 1 | 1 | 13 | 0 | 9 |

1 No estimate made.

City reports for week ended August 21, 1926—Continued

| Division, State, and city | Scarlet fever | | Smallpox | | | Tuber- culosis, deaths re- ported | Typhoid fever | | | Whoop- ing cough, cases re- ported | Deaths, all causes |
|------------------------------|---|------------------------|---|------------------------|-------------------------|---|---|------------------------|-------------------------|---|--------------------------|
| | Cases, esti- mated expect- ancy | Cases re- ported | Cases, esti- mated expect- ancy | Cases re- ported | Deaths re- ported | | Cases, esti- mated expect- ancy | Cases re- ported | Deaths re- ported | | |
| NEW ENGLAND | | | | | | | | | | | |
| Maine: | | | | | | | | | | | |
| Portland..... | 0 | 0 | 0 | 0 | 0 | 2 | 1 | 1 | 0 | 4 | 22 |
| New Hampshire: | | | | | | | | | | | |
| Concord..... | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 6 |
| Manchester..... | 0 | 2 | 0 | 0 | 0 | 1 | 9 | 0 | 0 | 0 | 14 |
| Vermont: | | | | | | | | | | | |
| Barre..... | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 1 |
| Massachusetts: | | | | | | | | | | | |
| Boston..... | 13 | 18 | 0 | 0 | 0 | 10 | 3 | 3 | 0 | 21 | 177 |
| Fall River..... | 1 | 0 | 0 | 0 | 0 | 5 | 2 | 0 | 0 | 1 | 31 |
| Springfield..... | 1 | 2 | 0 | 0 | 0 | 1 | 1 | 0 | 0 | 2 | 23 |
| Worcester..... | 2 | 5 | 0 | 0 | 0 | 2 | 0 | 0 | 0 | 3 | 28 |
| Rhode Island: | | | | | | | | | | | |
| Pawtucket..... | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 10 |
| Providence..... | 2 | 1 | 0 | 0 | 0 | 3 | 2 | 1 | 0 | 4 | 63 |
| Connecticut: | | | | | | | | | | | |
| Bridgeport..... | 2 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 1 | 27 |
| Hartford..... | 1 | 2 | 0 | 0 | 0 | 2 | 1 | 1 | 0 | 4 | 36 |
| New Haven..... | 1 | 2 | 0 | 0 | 0 | 1 | 3 | 1 | 0 | 2 | 22 |
| MIDDLE ATLANTIC | | | | | | | | | | | |
| New York: | | | | | | | | | | | |
| Buffalo..... | 4 | 5 | 0 | 0 | 0 | 10 | 3 | 1 | 1 | 1 | 115 |
| New York..... | 23 | 38 | 0 | 2 | 0 | 183 | 42 | 50 | 2 | 79 | 1,070 |
| Rochester..... | 3 | 2 | 0 | 0 | 0 | 1 | 1 | 2 | 0 | 9 | 57 |
| Syracuse..... | 2 | 0 | 0 | 0 | 0 | 0 | 1 | 1 | 0 | 9 | 42 |
| New Jersey: | | | | | | | | | | | |
| Camden..... | 0 | 1 | 1 | 0 | 0 | 2 | 1 | 0 | 0 | 7 | 27 |
| Newark..... | 4 | 1 | 0 | 0 | 0 | 9 | 2 | 2 | 0 | 24 | 85 |
| Trenton..... | 0 | 0 | 9 | 0 | 0 | 0 | 1 | 2 | 0 | 5 | 30 |
| Pennsylvania: | | | | | | | | | | | |
| Philadelphia..... | 16 | 11 | 0 | 0 | 0 | 28 | 13 | 10 | 3 | 42 | 405 |
| Pittsburgh..... | 8 | 0 | 0 | 0 | 0 | 3 | 4 | 0 | 0 | 33 | 122 |
| Reading..... | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 0 | 0 | 22 | 13 |
| EAST NORTH CENTRAL | | | | | | | | | | | |
| Ohio: | | | | | | | | | | | |
| Cincinnati..... | 3 | 3 | 0 | 0 | 0 | 12 | 3 | 1 | 0 | 5 | 141 |
| Cleveland..... | 7 | 15 | 0 | 0 | 0 | 19 | 5 | 5 | 1 | 73 | 164 |
| Columbus..... | 1 | 3 | 0 | 0 | 0 | 1 | 3 | 1 | 0 | 1 | 62 |
| Toledo..... | 5 | 1 | 1 | 1 | 0 | 5 | 3 | 2 | 1 | 39 | 73 |
| Indiana: | | | | | | | | | | | |
| Fort Wayne..... | 0 | 0 | 0 | 0 | 0 | 1 | 1 | 0 | 1 | 0 | 19 |
| Indianapolis..... | 2 | 1 | 1 | 2 | 0 | 1 | 3 | 1 | 1 | 21 | 103 |
| South Bend..... | 1 | 2 | 0 | 0 | 0 | 0 | 1 | 0 | 0 | 2 | 12 |
| Terre Haute..... | 1 | 0 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 16 |
| Illinois: | | | | | | | | | | | |
| Chicago..... | 23 | 18 | 0 | 1 | 0 | 49 | 6 | 7 | 0 | 71 | 536 |
| Peoria..... | 1 | 0 | 0 | 0 | 0 | 1 | 1 | 0 | 0 | 1 | 17 |
| Springfield..... | 0 | 1 | 0 | 0 | 0 | 1 | 1 | 2 | 0 | 3 | 13 |
| Michigan: | | | | | | | | | | | |
| Detroit..... | 20 | 17 | 2 | 0 | 0 | 15 | 5 | 8 | 2 | 80 | 201 |
| Flint..... | 3 | 2 | 0 | 0 | 0 | 2 | 1 | 0 | 0 | 2 | 17 |
| Grand Rapids..... | 1 | 1 | 0 | 0 | 0 | 2 | 1 | 0 | 0 | 5 | 34 |
| Wisconsin: | | | | | | | | | | | |
| Kenosha..... | 0 | 0 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 14 | 3 |
| Madison..... | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Milwaukee..... | 7 | 3 | 1 | 0 | 0 | 5 | 1 | 0 | 0 | 53 | 76 |
| Racine..... | 1 | 0 | 0 | 0 | 0 | 2 | 0 | 0 | 0 | 0 | 10 |
| Superior..... | 1 | 2 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 5 |
| WEST NORTH CENTRAL | | | | | | | | | | | |
| Minnesota: | | | | | | | | | | | |
| Duluth..... | 3 | 10 | 1 | 0 | 0 | 0 | 0 | 1 | 0 | 2 | 23 |
| Minneapolis..... | 10 | 17 | 1 | 0 | 0 | 4 | 1 | 0 | 0 | 2 | 73 |
| St. Paul..... | 4 | 5 | 1 | 0 | 0 | 4 | 3 | 5 | 0 | 11 | 52 |

1 Pulmonary tuberculosis only.

City reports for week ended August 21, 1926—Continued

| Division, State, and city | Scarlet fever | | Smallpox | | | Tuber- culosis, deaths re- ported | Typhoid fever | | | Whoop- ing cough, cases re- ported | Deaths, all causes |
|-----------------------------------|---|------------------------|---|------------------------|-------------------------|---|---|------------------------|-------------------------|---|--------------------------|
| | Cases, esti- mated expect- ancy | Cases re- ported | Cases, esti- mated expect- ancy | Cases re- ported | Deaths re- ported | | Cases, esti- mated expect- ancy | Cases re- ported | Deaths re- ported | | |
| WEST NORTH CEN- TRAL—continued | | | | | | | | | | | |
| Iowa: | | | | | | | | | | | |
| Davenport | 0 | 3 | 0 | 0 | | | 0 | 0 | | 1 | |
| Sioux City | 1 | 0 | 1 | 1 | | | 0 | 0 | | 9 | |
| Waterloo | 1 | 0 | 0 | 0 | | | 0 | 0 | | 4 | |
| Missouri: | | | | | | | | | | | |
| Kansas City | 2 | 0 | 0 | 0 | 0 | 4 | 3 | 0 | 0 | 3 | 93 |
| St. Joseph | 1 | 0 | 0 | 0 | 0 | 2 | 1 | 1 | 0 | 0 | 10 |
| St. Louis | 6 | 16 | 0 | 1 | 0 | 6 | 8 | 12 | 1 | 11 | 163 |
| North Dakota: | | | | | | | | | | | |
| Fargo | 1 | 4 | 0 | 0 | 0 | 1 | 0 | 0 | 0 | 0 | 14 |
| Grand Forks | 1 | 0 | 0 | 0 | | | 0 | 0 | | 0 | |
| South Dakota: | | | | | | | | | | | |
| Aberdeen | 0 | 0 | 0 | 0 | | | 0 | 0 | | 1 | |
| Sioux Falls | 1 | | 0 | | | | | | | | |
| Nebraska: | | | | | | | | | | | |
| Lincoln | 1 | 2 | 0 | 2 | 0 | 0 | 0 | 0 | 0 | 8 | 7 |
| Omaha | 1 | 2 | 1 | 0 | 0 | 1 | 1 | 1 | 1 | 3 | 47 |
| Kansas: | | | | | | | | | | | |
| Topeka | 1 | 4 | 0 | 0 | 0 | 0 | 1 | 3 | 0 | 5 | 20 |
| Wichita | 1 | 1 | 0 | 0 | 0 | 2 | 2 | 1 | 0 | 3 | 21 |
| SOUTH ATLANTIC | | | | | | | | | | | |
| Delaware: | | | | | | | | | | | |
| Wilmington | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 2 | 0 | 1 | 25 |
| Maryland: | | | | | | | | | | | |
| Baltimore | 5 | 5 | 0 | 0 | 0 | 20 | 9 | 11 | 2 | 72 | 212 |
| Cumberland | 1 | 0 | 0 | 0 | 0 | 0 | 1 | 0 | 0 | 0 | 10 |
| Frederick | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 3 |
| District of Col.: | | | | | | | | | | | |
| Washington | 2 | 6 | 0 | 0 | 0 | 6 | 5 | 4 | 0 | 14 | 90 |
| Virginia: | | | | | | | | | | | |
| Lynchburg | 1 | 1 | 0 | 0 | 0 | 0 | 1 | 2 | 0 | 4 | 16 |
| Norfolk | 0 | 1 | 0 | 1 | 0 | 2 | 2 | 4 | 0 | 3 | |
| Richmond | 2 | 5 | 0 | 0 | 0 | 1 | 2 | 6 | 1 | 1 | 31 |
| Roanoke | 0 | 0 | 0 | 0 | 0 | 1 | 2 | 1 | 0 | 0 | 20 |
| West Virginia: | | | | | | | | | | | |
| Charleston | 0 | 0 | 0 | 0 | 0 | 0 | 2 | 2 | 0 | 6 | 18 |
| Huntington | 0 | 0 | 0 | 0 | 0 | 1 | 2 | 0 | 0 | 0 | 25 |
| Wheeling | 1 | 0 | 0 | 0 | 0 | 0 | 1 | 0 | 1 | 1 | 16 |
| North Carolina: | | | | | | | | | | | |
| Raleigh | 1 | 1 | 0 | 0 | 0 | 1 | 0 | 0 | 0 | 26 | 16 |
| Wilmington | 1 | 0 | 0 | 0 | 0 | 1 | 1 | 1 | 0 | 3 | 9 |
| Winston-Salem | 0 | 0 | 1 | 0 | 0 | 1 | 3 | 0 | 0 | 0 | 21 |
| South Carolina: | | | | | | | | | | | |
| Charleston | 0 | 0 | 0 | 0 | 0 | 2 | 3 | 2 | 0 | 2 | 27 |
| Columbia | 1 | 1 | 0 | 0 | 0 | 0 | 1 | 1 | 0 | 0 | |
| Greenville | 0 | 0 | 1 | 0 | 0 | 0 | 1 | 0 | 0 | 0 | 8 |
| Georgia: | | | | | | | | | | | |
| Atlanta | 3 | 0 | 1 | 1 | 0 | 4 | 4 | 13 | 3 | 1 | 78 |
| Brunswick | 0 | 0 | 0 | 0 | 0 | 1 | 1 | 0 | 0 | 0 | 6 |
| Savannah | 1 | 0 | 0 | 0 | 0 | 2 | 2 | 0 | 0 | 0 | 30 |
| Florida: | | | | | | | | | | | |
| Miami | | 0 | | 0 | 0 | 0 | | 0 | 0 | 1 | 20 |
| St. Petersburg | 0 | | 0 | | 0 | 0 | 0 | | 0 | | 2 |
| Tampa | 0 | 1 | 0 | 1 | 0 | 3 | 0 | 1 | 2 | 0 | 34 |
| EAST SOUTH CEN- TRAL | | | | | | | | | | | |
| Kentucky: | | | | | | | | | | | |
| Covington | 0 | 0 | 1 | 0 | 0 | 1 | 0 | 1 | 0 | 0 | 19 |
| Louisville | 1 | 2 | 1 | 0 | 0 | 10 | 5 | 5 | 0 | 0 | 91 |
| Tennessee: | | | | | | | | | | | |
| Memphis | 1 | 0 | 0 | 0 | 0 | 3 | 7 | 16 | 2 | 8 | 49 |
| Nashville | 1 | 3 | 0 | 0 | 0 | 3 | 7 | 8 | 1 | 20 | 39 |
| Alabama: | | | | | | | | | | | |
| Birmingham | 3 | 2 | 0 | 0 | 0 | 5 | 7 | 2 | 0 | 17 | 57 |
| Mobile | 0 | 0 | 0 | 0 | 0 | 1 | 1 | 1 | 0 | 0 | 24 |
| Montgomery | 1 | 0 | 0 | 1 | 0 | 0 | 1 | 3 | 0 | 1 | 7 |

City reports for week ended August 21, 1928—Continued

| Division, State, and city | Scarlet fever | | Smallpox | | | Tuber- culosis, deaths re- ported | Typhoid fever | | | Whoop- ing cough, cases re- ported | Deaths, all causes |
|------------------------------|---|------------------------|---|------------------------|-------------------------|---|---|------------------------|-------------------------|---|--------------------------|
| | Cases, esti- mated expect- ancy | Cases re- ported | Cases, esti- mated expect- ancy | Cases re- ported | Deaths re- ported | | Cases, esti- mated expect- ancy | Cases re- ported | Deaths re- ported | | |
| WEST SOUTH CENTRAL | | | | | | | | | | | |
| Arkansas: | | | | | | | | | | | |
| Fort Smith..... | 1 | | 0 | | | | 0 | | | | |
| Little Rock..... | 0 | 0 | 0 | 0 | 0 | 3 | 2 | 0 | 0 | 0 | |
| Louisiana: | | | | | | | | | | | |
| New Orleans..... | 1 | 2 | 0 | 0 | 0 | 9 | 5 | 4 | 0 | 2 | 146 |
| Shreveport..... | 0 | 0 | 0 | 0 | 0 | 1 | 5 | 1 | 0 | 0 | 29 |
| Oklahoma: | | | | | | | | | | | |
| Oklahoma City..... | 1 | 1 | 0 | 0 | 0 | 0 | 2 | 2 | 0 | 0 | 23 |
| Texas: | | | | | | | | | | | |
| Dallas..... | 2 | 2 | 0 | 0 | 0 | 3 | 4 | 4 | 0 | 9 | 54 |
| Galveston..... | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 14 |
| Houston..... | 0 | 0 | 0 | 0 | 0 | 2 | 0 | 0 | 0 | 0 | 46 |
| San Antonio..... | 0 | 0 | 0 | 0 | 0 | 8 | 1 | 1 | 0 | 0 | 60 |
| MOUNTAIN | | | | | | | | | | | |
| Montana: | | | | | | | | | | | |
| Billings..... | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 6 |
| Great Falls..... | 1 | 0 | 0 | 0 | 0 | 0 | 1 | 0 | 0 | 0 | 3 |
| Helena..... | 0 | 0 | 0 | 0 | 0 | 1 | 0 | 0 | 0 | 0 | 5 |
| Missoula..... | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 1 | 0 | 0 | 4 |
| Idaho: | | | | | | | | | | | |
| Boise..... | 0 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 3 |
| Colorado: | | | | | | | | | | | |
| Denver..... | 2 | 3 | 2 | 0 | 0 | 4 | 3 | 1 | 0 | 3 | 65 |
| Pueblo..... | 0 | 0 | 0 | 0 | 0 | 2 | 1 | 5 | 0 | 0 | 9 |
| New Mexico: | | | | | | | | | | | |
| Albuquerque..... | 0 | 1 | 0 | 0 | 0 | 4 | 1 | 0 | 0 | 4 | 21 |
| Arizona: | | | | | | | | | | | |
| Phoenix..... | | 1 | 0 | 0 | 0 | 4 | 0 | 0 | 0 | 0 | 12 |
| Utah: | | | | | | | | | | | |
| Salt Lake City..... | 1 | 0 | 0 | 0 | 0 | 0 | 2 | 1 | 0 | 10 | 31 |
| Nevada: | | | | | | | | | | | |
| Reno..... | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 0 | 0 | 0 | 2 |
| PACIFIC | | | | | | | | | | | |
| Washington: | | | | | | | | | | | |
| Seattle..... | 3 | 1 | 1 | 1 | | | 2 | 2 | | 11 | |
| Spokane..... | 2 | 9 | 1 | 0 | | | 1 | 2 | | 9 | |
| Tacoma..... | 0 | 2 | 1 | 0 | 0 | 2 | 0 | 0 | 0 | 1 | 29 |
| Oregon: | | | | | | | | | | | |
| Portland..... | 3 | 9 | 4 | 1 | 0 | 3 | 1 | 1 | 0 | 1 | |
| California: | | | | | | | | | | | |
| Los Angeles..... | 6 | 9 | 2 | 1 | 0 | 20 | 4 | 3 | 0 | 4 | 196 |
| Sacramento..... | 1 | 0 | 1 | 0 | 0 | 0 | 1 | 1 | 1 | 0 | 29 |
| San Francisco..... | 5 | 8 | 0 | 0 | 0 | 6 | 2 | 1 | 0 | 2 | 150 |

| Division, State, and city | Cerebrospinal meningitis | | Lethargic encephalitis | | Pellagra | | Poliomyelitis (infantile paralysis) | |
|---------------------------|--------------------------|--------|------------------------|--------|----------|--------|-------------------------------------|--------|
| | Cases | Deaths | Cases | Deaths | Cases | Deaths | Cases, estimated expectancy | Deaths |
| NEW ENGLAND | | | | | | | | |
| Massachusetts: | | | | | | | | |
| Boston..... | 0 | 0 | 1 | 2 | 0 | 0 | 1 | 1 |
| Fall River..... | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 1 |
| Springfield..... | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 4 |
| Worcester..... | 0 | 0 | 1 | 1 | 0 | 0 | 0 | 10 |
| Rhode Island: | | | | | | | | |
| Providence..... | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 2 |
| MIDDLE ATLANTIC | | | | | | | | |
| New York: | | | | | | | | |
| Buffalo..... | 0 | 0 | 1 | 0 | 0 | 0 | 1 | 13 |
| New York..... | 3 | 4 | 5 | 4 | 0 | 0 | 8 | 5 |
| Syracuse..... | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 4 |

City reports for week ended August 21, 1926—Continued

| Division, State, and city | Cerebrospinal meningitis | | Lethargic encephalitis | | Pellagra | | Poliomyelitis (infantile paralysis) | | |
|---------------------------|--------------------------|--------|------------------------|--------|----------|--------|-------------------------------------|-------|--------|
| | Cases | Deaths | Cases | Deaths | Cases | Deaths | Cases, estimated expectancy | Cases | Deaths |
| EAST NORTH CENTRAL | | | | | | | | | |
| Ohio: | | | | | | | | | |
| Cleveland..... | 0 | 0 | 0 | 1 | 0 | 1 | 0 | 0 | 1 |
| Illinois: | | | | | | | | | |
| Chicago..... | 1 | 0 | 2 | 0 | 0 | 0 | 5 | 1 | 0 |
| Michigan: | | | | | | | | | |
| Flint..... | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 0 |
| WEST NORTH CENTRAL | | | | | | | | | |
| Missouri: | | | | | | | | | |
| Kansas City..... | 0 | 0 | 0 | 0 | 1 | 1 | 0 | 0 | 0 |
| St. Louis..... | 1 | 0 | 0 | 0 | 0 | 0 | 1 | 0 | 0 |
| SOUTH ATLANTIC | | | | | | | | | |
| Maryland: | | | | | | | | | |
| Baltimore..... | 0 | 0 | 2 | 0 | 0 | 0 | 1 | 2 | 2 |
| Virginia: | | | | | | | | | |
| Roanoke..... | 0 | 0 | 0 | 0 | 0 | 1 | 0 | 0 | 0 |
| North Carolina: | | | | | | | | | |
| Raleigh..... | 0 | 0 | 0 | 0 | 0 | 1 | 0 | 0 | 0 |
| Winston-Salem..... | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| South Carolina: | | | | | | | | | |
| Charleston..... | 0 | 0 | 1 | 0 | 1 | 0 | 0 | 0 | 0 |
| Georgia: | | | | | | | | | |
| Brunswick..... | 0 | 0 | 0 | 0 | 0 | 1 | 0 | 0 | 0 |
| Florida: | | | | | | | | | |
| St. Petersburg..... | 0 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| EAST SOUTH CENTRAL | | | | | | | | | |
| Kentucky: | | | | | | | | | |
| Louisville..... | 0 | 0 | 1 | 2 | 0 | 0 | 0 | 0 | 0 |
| WEST SOUTH CENTRAL | | | | | | | | | |
| Arkansas: | | | | | | | | | |
| Little Rock..... | 0 | 0 | 0 | 0 | 0 | 2 | 0 | 0 | 0 |
| Louisiana: | | | | | | | | | |
| New Orleans..... | 0 | 0 | 0 | 0 | 2 | 2 | 0 | 0 | 0 |
| Oklahoma: | | | | | | | | | |
| Oklahoma City..... | 0 | 0 | 0 | 0 | 1 | 0 | 0 | 0 | 0 |
| Texas: | | | | | | | | | |
| Galveston..... | 0 | 0 | 0 | 0 | 0 | 2 | 0 | 0 | 0 |
| MOUNTAIN | | | | | | | | | |
| Colorado: | | | | | | | | | |
| Denver..... | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 2 | 0 |
| Utah: | | | | | | | | | |
| Salt Lake City..... | 0 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| PACIFIC | | | | | | | | | |
| Washington: | | | | | | | | | |
| Spokane..... | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 0 |
| California: | | | | | | | | | |
| Los Angeles..... | 0 | 0 | 1 | 0 | 0 | 0 | 0 | 1 | 0 |
| Sacramento..... | 1 | 1 | 0 | 0 | 0 | 0 | 1 | 0 | 0 |
| San Francisco..... | 0 | 0 | 0 | 0 | 0 | 1 | 0 | 0 | 0 |

The following table gives the rates per 100,000 population for 102 cities for the five-week period ended August 21, 1926, compared with those for a like period ended August 22, 1925. The population figures used in computing the rates are approximate estimates as of July 1, 1925 and 1926, respectively, authoritative figures for many of the cities not being available. The 102 cities reporting cases had an

estimated aggregate population of nearly 30,000,000 in 1925 and nearly 30,500,000 in 1926. The 96 cities reporting deaths had more than 29,250,000 estimated population in 1925 and more than 29,750,000 in 1926. The number of cities included in each group and the estimated aggregate populations are shown in a separate table below.

Summary of weekly reports from cities, July 18 to August 21, 1926—Annual rates per 100,000 population—Compared with rates for the corresponding period of 1925¹

DIPHTHERIA CASE RATES

| | Week ended— | | | | | | | | | |
|-------------------------|---------------|-----------------|-----------------|-----------------|------------------|------------------|---------------|------------------|---------------|------------------|
| | July 25, 1925 | July 24, 1926 | Aug. 1, 1925 | July 31, 1926 | Aug. 8, 1925 | Aug. 7, 1926 | Aug. 15, 1925 | Aug. 14, 1926 | Aug. 22, 1925 | Aug. 21, 1926 |
| 102 cities..... | 75 | ² 90 | ³ 75 | ² 80 | ⁴ 83 | ³ 78 | 77 | ⁵ 69 | 68 | ⁶ 63 |
| New England..... | 60 | 33 | 60 | 40 | 79 | 40 | 89 | 31 | 50 | 47 |
| Middle Atlantic..... | 90 | 109 | 92 | 103 | 83 | 88 | 78 | 62 | 73 | 59 |
| East North Central..... | 63 | 99 | 69 | 83 | 94 | ⁷ 105 | 68 | ⁷ 101 | 51 | ⁷ 87 |
| West North Central..... | 103 | ² 95 | 97 | ² 85 | ¹ 105 | ² 82 | 107 | ¹ 56 | 99 | ³ 85 |
| South Atlantic..... | 42 | 34 | ⁴ 48 | 21 | 52 | 43 | 69 | 49 | 60 | 60 |
| East South Central..... | 11 | 10 | 11 | 21 | 26 | 10 | 32 | 57 | 58 | 21 |
| West South Central..... | 66 | 39 | 40 | 39 | 22 | 39 | 48 | 26 | 57 | ¹⁰ 66 |
| Mountain..... | 111 | 64 | 148 | 91 | ¹¹ 66 | 118 | 157 | 73 | 74 | 146 |
| Pacific..... | 99 | 175 | 64 | 119 | 141 | 102 | 80 | 105 | 110 | 62 |

MEASLES CASE RATES

| 102 cities..... | 101 | ² 155 | ³ 70 | ² 103 | ⁴ 51 | ⁵ 66 | 46 | ⁶ 57 | 30 | ⁷ 41 |
|-------------------------|-----|------------------|-----------------|------------------|------------------|-----------------|-----|-----------------|----|-----------------|
| New England..... | 208 | 109 | 180 | 83 | 127 | 83 | 125 | 69 | 93 | 52 |
| Middle Atlantic..... | 127 | 108 | 77 | 63 | 69 | 42 | 57 | 33 | 88 | 27 |
| East North Central..... | 111 | 243 | 68 | 171 | 44 | ⁷ 96 | 35 | ⁷ 77 | 21 | ⁷ 60 |
| West North Central..... | 18 | ¹ 183 | 30 | ³ 93 | ⁸ 10 | ³ 58 | 24 | ² 66 | 6 | ⁹ 29 |
| South Atlantic..... | 90 | 128 | ³ 68 | 115 | 42 | 47 | 40 | 81 | 33 | 36 |
| East South Central..... | 58 | 125 | 26 | 93 | 11 | 42 | 16 | 31 | 5 | 36 |
| West South Central..... | 4 | 13 | 0 | 9 | 0 | 9 | 9 | 4 | 9 | ¹⁰ 9 |
| Mountain..... | 37 | 173 | 102 | 127 | ¹¹ 19 | 137 | 18 | 64 | 28 | 18 |
| Pacific..... | 19 | 213 | 33 | 121 | 28 | 121 | 19 | 94 | 11 | 78 |

SCARLET FEVER CASE RATES

| 102 cities..... | 55 | ² 83 | ³ 54 | ⁴ 73 | ⁵ 51 | ⁶ 61 | 57 | ⁷ 51 | 51 | ⁸ 48 |
|-------------------------|-----|------------------|-----------------|------------------|------------------|------------------|-----|------------------|-----|-------------------|
| New England..... | 69 | 85 | 72 | 118 | 98 | 104 | 81 | 69 | 80 | 73 |
| Middle Atlantic..... | 42 | 75 | 37 | 52 | 33 | 38 | 36 | 30 | 23 | 29 |
| East North Central..... | 63 | 93 | 60 | 85 | 48 | ⁷ 79 | 54 | ⁷ 66 | 54 | ⁷ 47 |
| West North Central..... | 115 | ² 127 | 121 | ² 143 | ⁸ 117 | ¹ 101 | 129 | ² 119 | 143 | ¹² 123 |
| South Atlantic..... | 15 | 36 | ³ 34 | 34 | 21 | 39 | 38 | 30 | 40 | 39 |
| East South Central..... | 26 | 93 | 58 | 62 | 58 | 31 | 37 | 47 | 32 | 36 |
| West South Central..... | 31 | 82 | 26 | 39 | 53 | 17 | 66 | 22 | 48 | ¹⁰ 18 |
| Mountain..... | 157 | 64 | 83 | 36 | ¹¹ 38 | 64 | 92 | 36 | 65 | 36 |
| Pacific..... | 44 | 92 | 47 | 86 | 61 | 84 | 83 | 86 | 41 | 78 |

¹ The figures given in this table are rates per 100,000 population, annual basis—and not the number of cases reported. Populations used are estimated as of July 1, 1925 and 1926, respectively.

² Sioux Falls, S. Dak., not included.

³ Tampa, Fla., not included.

⁴ Waterloo, Iowa, and Helena, Mont., not included.

⁵ Madison, Wis., and Sioux Falls, S. Dak., not included.

⁶ Madison, Wis., Sioux City, Iowa, Sioux Falls, S. Dak., and Fort Smith, Ark., not included.

⁷ Madison, Wis., not included.

⁸ Waterloo, Iowa, not included.

⁹ Sioux City, Iowa, and Sioux Falls, S. Dak., not included.

¹⁰ Fort Smith, Ark., not included.

¹¹ Helena, Mont., not included.

Summary of weekly reports from cities, July 18 to August 21, 1926—Annual rates per 100,000 population—Compared with rates for the corresponding period of 1925—Continued

SMALLPOX CASE RATES

| | Week ended— | | | | | | | | | |
|-------------------------|---------------|---------------|--------------|---------------|--------------|--------------|---------------|---------------|---------------|---------------|
| | July 25, 1925 | July 24, 1926 | Aug. 1, 1925 | July 31, 1926 | Aug. 8, 1925 | Aug. 7, 1926 | Aug. 15, 1925 | Aug. 14, 1926 | Aug. 22, 1925 | Aug. 21, 1926 |
| 102 cities..... | 10 | 16 | 19 | 15 | 19 | 18 | 7 | 17 | 6 | 12 |
| New England..... | 5 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Middle Atlantic..... | 0 | 0 | 0 | 1 | 0 | 1 | 0 | 0 | 0 | 1 |
| East North Central..... | 8 | 8 | 3 | 6 | 6 | 79 | 3 | 11 | 2 | 12 |
| West North Central..... | 12 | 14 | 14 | 14 | 18 | 14 | 16 | 14 | 6 | 12 |
| South Atlantic..... | 15 | 6 | 12 | 2 | 2 | 11 | 2 | 11 | 4 | 6 |
| East South Central..... | 37 | 10 | 21 | 5 | 47 | 16 | 21 | 26 | 37 | 5 |
| West South Central..... | 4 | 13 | 4 | 4 | 13 | 13 | 9 | 22 | 4 | 10 |
| Mountain..... | 0 | 27 | 55 | 9 | 119 | 9 | 9 | 73 | 9 | 0 |
| Pacific..... | 64 | 8 | 80 | 32 | 64 | 24 | 64 | 32 | 41 | 5 |

TYPHOID FEVER CASE RATES

| 102 cities..... | 33 | 18 | 40 | 30 | 40 | 29 | 46 | 35 | 55 | 41 |
|-------------------------|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| New England..... | 22 | 9 | 22 | 14 | 26 | 12 | 38 | 17 | 31 | 17 |
| Middle Atlantic..... | 21 | 9 | 30 | 23 | 23 | 19 | 33 | 24 | 44 | 34 |
| East North Central..... | 8 | 6 | 10 | 10 | 20 | 12 | 17 | 19 | 29 | 17 |
| West North Central..... | 38 | 12 | 46 | 22 | 41 | 18 | 35 | 24 | 46 | 56 |
| South Atlantic..... | 50 | 47 | 64 | 54 | 56 | 66 | 86 | 100 | 104 | 94 |
| East South Central..... | 163 | 135 | 168 | 259 | 252 | 182 | 200 | 140 | 168 | 187 |
| West South Central..... | 163 | 30 | 154 | 47 | 123 | 60 | 97 | 47 | 128 | 44 |
| Mountain..... | 46 | 46 | 55 | 36 | 104 | 27 | 102 | 73 | 102 | 73 |
| Pacific..... | 28 | 8 | 44 | 11 | 17 | 30 | 41 | 30 | 61 | 24 |

INFLUENZA DEATH RATES

| 96 cities..... | 2 | 3 | 1 | 2 | 12 | 2 | 2 | 1 | 2 | 3 |
|-------------------------|---|----|----|----|----|----|---|----|----|----|
| New England..... | 0 | 2 | 0 | 0 | 5 | 0 | 0 | 0 | 0 | 0 |
| Middle Atlantic..... | 3 | 2 | 1 | 1 | 2 | 2 | 3 | 1 | 2 | 1 |
| East North Central..... | 1 | 4 | 0 | 1 | 3 | 1 | 3 | 10 | 1 | 13 |
| West North Central..... | 4 | 12 | 0 | 10 | 0 | 10 | 0 | 12 | 0 | 12 |
| South Atlantic..... | 4 | 4 | 12 | 2 | 6 | 4 | 0 | 0 | 0 | 2 |
| East South Central..... | 5 | 5 | 0 | 5 | 5 | 0 | 5 | 10 | 11 | 0 |
| West South Central..... | 0 | 9 | 0 | 24 | 5 | 5 | 0 | 14 | 10 | 28 |
| Mountain..... | 9 | 9 | 0 | 0 | 10 | 9 | 9 | 0 | 9 | 0 |
| Pacific..... | 0 | 4 | 0 | 4 | 0 | 11 | 0 | 0 | 7 | 7 |

PNEUMONIA DEATH RATES

| 96 cities..... | 48 | 54 | 59 | 48 | 52 | 54 | 60 | 50 | 53 | 54 |
|-------------------------|----|----|-----|----|----|-----|----|-----|----|----|
| New England..... | 50 | 33 | 53 | 33 | 36 | 54 | 29 | 31 | 38 | 40 |
| Middle Atlantic..... | 51 | 64 | 65 | 41 | 65 | 56 | 73 | 62 | 65 | 58 |
| East North Central..... | 37 | 46 | 48 | 48 | 36 | 42 | 47 | 35 | 40 | 34 |
| West North Central..... | 40 | 40 | 40 | 57 | 51 | 51 | 42 | 25 | 30 | 49 |
| South Atlantic..... | 52 | 58 | 60 | 51 | 50 | 68 | 73 | 56 | 60 | 86 |
| East South Central..... | 58 | 60 | 68 | 62 | 63 | 52 | 58 | 52 | 74 | 36 |
| West South Central..... | 63 | 57 | 116 | 76 | 68 | 104 | 82 | 113 | 77 | 71 |
| Mountain..... | 55 | 64 | 74 | 55 | 28 | 64 | 55 | 82 | 65 | 82 |
| Pacific..... | 58 | 35 | 62 | 71 | 69 | 57 | 80 | 39 | 47 | 78 |

¹ Sioux Falls, S. Dak., not included.

² Tampa, Fla., not included.

³ Waterloo, Iowa, and Helena, Mont., not included.

⁴ Madison, Wis., and Sioux Falls, S. Dak., not included.

⁵ Madison, Wis., Sioux City, Iowa, Sioux Falls, S. Dak., and Fort Smith, Ark., not included.

⁶ Madison, Wis., not included.

⁷ Waterloo, Iowa, not included.

⁸ Sioux City, Iowa, and Sioux Falls, S. Dak., not included.

⁹ Fort Smith, Ark., not included.

¹⁰ Helena, Mont., not included.

Number of cities included in summary of weekly reports, and aggregate population of cities in each group, approximated as of July 1, 1925 and 1926, respectively

| Group of cities | Number of cities reporting cases | Number of cities reporting deaths | Aggregate population of cities reporting cases | | Aggregate population of cities reporting deaths | |
|-------------------------|----------------------------------|-----------------------------------|--|------------|---|------------|
| | | | 1925 | 1926 | 1925 | 1926 |
| Total..... | 102 | 96 | 20,930,185 | 30,458,186 | 20,251,658 | 29,764,201 |
| New England..... | 12 | 12 | 2,176,124 | 2,206,124 | 2,176,124 | 2,206,124 |
| Middle Atlantic..... | 10 | 10 | 10,346,970 | 10,476,970 | 10,346,970 | 10,476,970 |
| East North Central..... | 16 | 16 | 7,481,656 | 7,653,436 | 7,481,656 | 7,655,436 |
| West North Central..... | 13 | 11 | 2,580,151 | 2,619,719 | 2,461,380 | 2,499,036 |
| South Atlantic..... | 21 | 21 | 2,716,070 | 2,776,070 | 2,716,070 | 2,776,070 |
| East South Central..... | 7 | 7 | 993,103 | 1,004,933 | 993,103 | 1,004,933 |
| West South Central..... | 8 | 6 | 1,184,057 | 1,212,057 | 1,078,198 | 1,103,695 |
| Mountain..... | 9 | 9 | 563,912 | 572,773 | 563,912 | 572,773 |
| Pacific..... | 6 | 4 | 1,888,142 | 1,934,084 | 1,434,245 | 1,469,144 |

FOREIGN AND INSULAR

CHOLERA ON VESSEL

Steamship "Macedonia"—Yokohama, Japan—August 5, 1926.—On August 5, 1926, a case of cholera was found on the steamship *Macedonia* at Yokohama, Japan. The *Macedonia* sailed from Singapore July 18, 1926.

THE FAR EAST

Report for week ended August 7, 1926.—The following report for the week ended August 7, 1926, was transmitted by the far eastern bureau of the health section of the secretariat of the League of Nations to the headquarters at Geneva:

| Maritime towns | Plague | | Cholera | | Small-pox | | Maritime towns | Plague | | Cholera | | Small-pox | |
|-----------------------|--------|--------|---------|--------|-----------|--------|-----------------------------|--------|--------|---------|--------|-----------|--------|
| | Cases | Deaths | Cases | Deaths | Cases | Deaths | | Cases | Deaths | Cases | Deaths | Cases | Deaths |
| Egypt—Alexandria..... | 0 | 1 | 0 | 0 | 2 | 1 | Siam—Bangkok..... | 0 | 0 | 8 | 2 | 4 | 4 |
| British India: | | | | | | | Dutch East Indies— | | | | | | |
| Bombay..... | 0 | 0 | 0 | 0 | 7 | 3 | Cheribon ¹ | 0 | 0 | 0 | 0 | 1 | 0 |
| Madras..... | 0 | 0 | 1 | 12 | 0 | 0 | French Indo-China— | | | | | | |
| Rangoon..... | 5 | 1 | 0 | 0 | 0 | 0 | Saigon and Cholon... | 1 | 0 | 1 | 0 | 0 | 0 |
| Negapatam..... | 0 | 0 | 3 | 0 | 0 | 0 | China: | | | | | | |
| Karachi..... | 0 | 0 | 0 | 1 | 0 | 0 | Amoy..... | 2 | 0 | 0 | 0 | 0 | 0 |
| Vizagapatam..... | 0 | 0 | 0 | 2 | 0 | 0 | Shanghai..... | | | | | | |

¹ One plague-infected rat was found in the port during the week.

Telegraphic reports from the following maritime towns indicated that no case of plague, cholera, or smallpox was reported during the week:

ASIA

Iraq.—Basra.

British India.—Chittagong, Cochin, Tuticorin.

Ceylon.—Colombo.

Federated Malay States.—Port Swettenham.

Straits Settlements.—Penang, Singapore.

Dutch East Indies.—Batavia, Surabaya, Samarang, Belawan-Deli, Palembang, Sabang, Makassar, Menado, Banjarmasin, Balikpapan, Tarakan, Padang, Samarinda.

Sarawak.—Kuching.

British North Borneo.—Sandakan, Jesselton, Kudat, Tawao.

Portuguese Timor.—Dilly.

Philippine Islands.—Manila, Iloilo, Jolo, Cebu, Zamboanga.

French Indo-China.—Turane, Haiphong.

China.—Hongkong.

Formosa.—Keelung.

Kwantung.—Port Arthur, Dairen.

Japan.—Yokohama, Osaka, Nagasaki, Moji, Kobe, Niigata, Tsuruga, Hakodate, Simonoseki.

Korea.—Chemulpo, Fusan.

Manchuria.—Antung, Mukden, Changchun, Harbin.

U. S. S. R.—Vladivostok.

AUSTRALASIA AND OCEANIA

Australia.—Adelaide, Melbourne, Sydney, Brisbane, Rockhampton, Townsville, Port Darwin, Broome, Fremantle, Carnarvon, Thursday Island.

New Guinea.—Port Moresby.

New Zealand.—Auckland, Wellington, Christchurch, Invercargill, Dunedin.

New Caledonia.—Noumea.

Fiji.—Suva.

Hawaii.—Honolulu.

AFRICA

Egypt.—Port Said, Suez.

Anglo-Egyptian Sudan.—Port Sudan, Suakin.

Eritrea.—Massaua.

French Somaliland.—Jibuti.

British Somaliland.—Berbera.

Italian Somaliland.—Mogadiscio.

Kenya.—Mombasa.

Zanzibar.—Zanzibar.

Tanganyika.—Dar-es-Salaam.

Seychelles.—Victoria.

Mauritius.—Port Louis.

Madagascar.—Tamatave, Majunga.

Portuguese East Africa.—Mozambique, Beira, Lourenço-Marques.

Union of South Africa.—Durban, East London, Port Elizabeth, Cape Town.

Reports had not been received in time for distribution from—

British India.—Calcutta.

Dutch East Indies.—Pontianak.

China.—Shanghai.

ALGERIA

Plague—Bona—August 14, 1926.—A case of plague was reported, August 14, 1926, at Bona, Algeria.

CANADA

Communicable diseases—Week ended August 21, 1926.—The Canadian Ministry of Health reports cases of certain communicable diseases in six Provinces of Canada for the week ended August 21, 1926, as follows:

| Disease | Nova Scotia | New Brunswick | Quebec | Ontario | Manitoba | Saskatchewan | Total |
|--------------------------|-------------|---------------|--------|---------|----------|--------------|-------|
| Cerebrospinal fever..... | | | 1 | 1 | | 1 | 3 |
| Influenza..... | 10 | | | | | | 10 |
| Polio-myelitis..... | | 2 | | 2 | | | 4 |
| Smallpox..... | | | 7 | 7 | 2 | 1 | 10 |
| Typhoid fever..... | 2 | 4 | 7 | 9 | 6 | | 28 |

Vital statistics—Quebec—June, 1926.—Births and deaths in the Province of Quebec for the month of June, 1926, have been reported as follows:

| | | | |
|--------------------------------------|-----------|--|-----|
| Estimated population..... | 2,570,000 | Deaths from—Continued. | |
| Births..... | 6,653 | Heart diseases..... | 404 |
| Birth rate per 1,000 population..... | 31.06 | Influenza..... | 95 |
| Deaths (all causes)..... | 2,884 | Measles..... | 30 |
| Death rate per 1,000 population..... | 13.46 | Poliomyelitis (infantile paralysis)..... | 1 |
| Deaths under 1 year..... | 837 | Scarlet fever..... | 14 |
| Infant mortality rate..... | 125.80 | Syphilis..... | 18 |
| Deaths from— | | Tuberculosis (pulmonary)..... | 235 |
| Cancer..... | 123 | Tuberculosis (other forms)..... | 69 |
| Cerebrospinal meningitis..... | 10 | Typhoid fever..... | 22 |
| Diabetes..... | 20 | Whooping cough..... | 41 |
| Diphtheria..... | 32 | | |

CANARY ISLANDS

Plague—Teneriffe—August 2, 1926.—Information received under date of August 5, 1926, shows two cases of plague reported present August 2, 1926, at Teneriffe, Canary Islands, occurring in the Cristianos district.

CHINA

Cholera—Swatow.—During the week ended July 31, 1926, 14 cases of cholera were reported at Swatow, China. It was stated that the mortality from the disease was low.

ECUADOR

Plague—Guayaquil—July 16–31, 1926.—During the period July 16 to 31, 1926, five cases of plague with two deaths were reported at Guayaquil, Ecuador.

Plague-infected rats.—During the same period, of 10,148 rats taken 14 were found plague infected.

EGYPT

Plague—July 23–29, 1926.—Plague has been reported in Egypt as follows: In cities—Alexandria, July 27, two cases; Suez, July 29, two cases, bubonic; in Provinces—Behera, July 23–29, two cases, bubonic; Charkieh, July 27, one case fatal, septicemic; Minieh, July 24, one case, fatal, bubonic.

Plague in Egypt—January 1–July 29, 1926

| Place | Cases | Deaths | Date of first case 1926 | Date of last case 1926 |
|-----------------|-------|--------|-------------------------|------------------------|
| City: | | | | |
| Alexandria..... | 5 | 1 | Mar. 10 | July 27 |
| Suez..... | 18 | 11 | Mar. 27 | July 29 |
| Province: | | | | |
| Behera..... | 20 | 7 | June 3 | July 29 |
| Beni-Suef..... | 45 | 24 | May 19 | July 15 |
| Charkieh..... | 1 | 1 | June 27 | July 27 |
| Dakahlia..... | 1 | 1 | Apr. 22 | May 22 |
| Fayoum..... | 1 | 1 | May 4 | May 4 |
| Gharbieh..... | 12 | 6 | Mar. 9 | June 2 |
| Girgeh..... | 6 | 4 | June 26 | July 3 |
| Minieh..... | 3 | 2 | Mar. 4 | July 24 |

FINLAND

Communicable diseases—June, 1926.—During the month of June, 1926, communicable diseases were reported in the Republic of Finland as follows:

| Disease | Cases | Disease | Cases |
|-----------------------------|-------|--------------------|-------|
| Diphtheria..... | 42 | Poliomyelitis..... | 3 |
| Dysentery..... | 6 | Scarlet fever..... | 96 |
| Lethargic encephalitis..... | 2 | Typhoid fever..... | 40 |
| Paratyphoid fever..... | 135 | | |

FRANCE

Plague—Saint Ouen—August 14, 1926.—Two cases of plague were reported at St. Ouen, a suburb of Paris, France, August 14, 1926.

GERMANY

Mortality—Karlsruhe, Baden—Year 1925 (comparative).—Information received under date of July 30, 1926, relative to vital statistics of the city of Karlsruhe, Baden, shows for the year 1925 the occurrence of 1,748 deaths from all causes in a population of 144,700; previous year, 1,738. For the years 1918, 1921, and 1923 the number of deaths were 2,320, 1,907, and 1,962, respectively. The decrease for the years 1924 and 1925 was stated to have been mainly due to decreased infant mortality, and this improvement was attributed to improved living conditions and better education in the care and treatment of infants.

Causes of death.—During the year 1925 tuberculosis caused 206 deaths, of which 167 were of tuberculosis of the lungs and larynx.

Cancer.—Increased mortality from cancer has been noted. In 1924 there were reported 206 deaths from cancer; in 1925 the number fell to 199; from 1920 to 1923 the number varied from 154 to 146.

GUATEMALA

Gastroenteritis—Guatemala—July, 1926.—During the month of July, 27 deaths from gastroenteritis were reported at Guatemala. Population, 220,000.

JAPAN

Further relative to plague at Yokohama.—Under date of August 7, 1926, two additional cases of plague¹ were reported at Yokohama, Japan, occurring within the city limits but not at the same distance from the original focus. One case occurred in a coolie employed in the canal section and one in an employee of a silk warehouse, a large reinforced building outside the customs compound.

MEXICO

Gastroenteritis—Chihuahua—Mazatlan—August 16–22, 1926.—During the week ended August 22, 1926, 7 deaths from gastroenteritis were reported at Chihuahua and 7 at Mazatlan, Mexico. Population, 48,000 and 25,000, respectively.

VENEZUELA

Gastroenteritis—Caracas—July, 1926.—During the month of July, 1926, 58 deaths from gastroenteritis were reported at Caracas, Venezuela. Of these, 42 deaths were in children under two years of age; 16 were in persons over two years.

VIRGIN ISLANDS

Communicable diseases—July, 1926.—Communicable diseases were reported in the Virgin Islands of the United States during the month of July, 1926, as follows:

| Island and disease | Cases | Remarks |
|--------------------------|-------|-------------------|
| St. Thomas and St. John: | | |
| Chancroid | 1 | St. Croix. |
| Gonorrhea | 4 | St. John, I. |
| Syphilis | 3 | Secondary. |
| Tuberculosis | 2 | Chronic pulmonary |
| St. Croix: | | |
| Chancroid | 6 | |
| Gonorrhea | 3 | |
| Leprosy | 1 | |
| Mumps | 2 | |

YUGOSLAVIA

Communicable diseases—July, 1926.—During the month of July, 1926, communicable diseases were reported in Yugoslavia as follows:

| Disease | Cases | Deaths | Disease | Cases | Deaths |
|--------------------------|-------|--------|----------------|-------|--------|
| Anthrax | 31 | 4 | Rabies | 4 | 4 |
| Cerebrospinal meningitis | 6 | 9 | Scarlet fever | 285 | 68 |
| Diphtheria | 97 | 8 | Tetanus | 30 | 19 |
| Dysentery | 166 | 18 | Typhoid fever | 198 | 23 |
| Glanders | 1 | 1 | Typhus fever | 2 | 1 |
| Measles | 277 | 6 | Whooping cough | 225 | 3 |

CHOLERA, PLAGUE, SMALLPOX, TYPHUS FEVER, AND YELLOW FEVER

The reports contained in the following tables must not be considered as complete or final as regards either the lists of countries included or the figures for the particular countries for which reports are given.

Reports Received During Week Ended September 10, 1926¹**CHOLERA**

| Place | Date | Cases | Deaths | Remarks |
|--------------------------|--------------------|-------|--------|---|
| China: | | | | |
| Shanghai..... | July 25-Aug. 1.... | 8 | 189 | Cases, foreign; deaths, native and foreign. |
| Swatow..... | July 18-24..... | | 48 | Reported by police. Possibly not all actually cholera cases. |
| Do..... | July 25-31..... | 14 | | Mortality stated to be low. |
| Tsingtao..... | do..... | | 1 | |
| India: | | | | |
| Bombay..... | July 18-24..... | 1 | 1 | |
| Indo-China: | | | | |
| Saigon..... | July 4-17..... | 8 | 2 | |
| On vessel: | | | | |
| Steamship Macedonia..... | Aug. 5..... | 1 | | At Yokohama, Japan. Vessel sailed from Singapore July 18, 1926. |

PLAGUE

| | | | | |
|------------------------|---------------------|----|----|--|
| Algeria: | | | | |
| Bona..... | Aug. 14..... | 1 | | |
| British East Africa: | | | | |
| Uganda..... | | | | Apr. 1-30, 1926: Cases, 100; deaths, 88. May, 1926: Cases, 314; deaths, 234. |
| Canary Islands: | | | | |
| Teneriffe..... | Aug. 2..... | 2 | | |
| China: | | | | |
| Foochow..... | July 11-24..... | | | Present, not epidemic. |
| Swatow..... | July 25-31..... | 14 | | |
| Ecuador: | | | | |
| Guayaquil..... | July 16-31..... | 5 | 2 | Rats taken, 10,146; found infected, 14. |
| Egypt: | | | | |
| City— | | | | Jan. 1-July 22, 1926: Cases, 104; for corresponding period, 1925— |
| Alexandria..... | July 27..... | 2 | | Cases, 88. |
| Suez..... | July 29..... | 2 | | Bubonic. |
| Province— | | | | |
| Behera..... | July 23-29..... | 2 | | Do. |
| Charkieh..... | July 27..... | 1 | 1 | Septicemic. |
| Minieh..... | July 24..... | 1 | 1 | Bubonic. |
| France: | | | | |
| Saint Ouen..... | Aug. 14..... | 2 | | Suburb of Paris. |
| Greece: | | | | |
| Patras..... | July 25-Aug. 7..... | 5 | 2 | |
| India: | | | | |
| Bombay..... | July 18-24..... | 1 | 1 | |
| Madras Presidency..... | July 4-10..... | 28 | 15 | |
| Iraq: | | | | |
| Baghdad..... | July 18-24..... | 1 | 1 | |
| Japan: | | | | |
| Yokohama..... | Aug. 7..... | 2 | | |
| Java: | | | | |
| Batavia..... | July 10-16..... | 7 | 7 | Province. |

SMALLPOX

| | | | | |
|---------------------|-----------------|-----|----|--|
| Belgium: | | | | |
| Antwerp..... | Aug. 1-7..... | 1 | 1 | |
| Brazil: | | | | |
| Bahia..... | July 11-17..... | 1 | | |
| Pernambuco..... | July 11-17..... | 1 | | |
| Rio de Janeiro..... | July 4-10..... | 54 | 22 | Jan. 1-July 31, 1926: Cases, 1,379; deaths, 689. |
| Do..... | July 11-17..... | 126 | 48 | |
| Do..... | July 18-24..... | 142 | 80 | |
| Do..... | July 25-31..... | 186 | 85 | |

¹ From medical officers of the Public Health Service, American consuls, and other sources.

CHOLERA, PLAGUE, SMALLPOX, TYPHUS FEVER, AND YELLOW FEVER—Continued**Reports Received During Week Ended September 10, 1926—Continued****SMALLPOX—Continued**

| Place | Date | Cases | Deaths | Remarks |
|----------------------------|---------------------|-------|--------|---|
| British East Africa: | | | | |
| Mombasa..... | July 5-11..... | 5 | 4 | May, 1926: Cases, 252; deaths, 46. April, 1926: One case. May, 1926: One case. |
| Tanganyika..... | | | | |
| Uganda..... | | | | |
| Canada: | | | | |
| British Columbia— | | | | |
| Vancouver..... | Aug. 16-22..... | 2 | | Aug. 15-21, 1926: Cases, 2. |
| Manitoba..... | | | | |
| Winnipeg..... | Aug. 15-23..... | 5 | | Aug. 15-21, 1926: Cases, 7. Aug. 15-21, 1926: Case, 1. |
| Ontario..... | | | | |
| Saskatchewan..... | | | | |
| China: | | | | |
| Chungking..... | July 18-31..... | | | Present. Do. |
| Foochow..... | July 11-24..... | | | |
| Manchuria— | | | | |
| Harbin..... | July 22-23..... | 5 | | At stations. Sporadic. |
| South Manchurian Ry. | July 25-31..... | 5 | | |
| Swatow..... | July 18-31..... | | | |
| Great Britain: | | | | |
| England and Wales..... | | | | Aug. 1-14, 1926: Cases, 134. |
| Sheffield..... | Aug. 1-7..... | 1 | | |
| India: | | | | |
| Bombay..... | July 18-24..... | 16 | 4 | |
| Madras..... | July 25-31..... | 3 | 1 | |
| Java: | | | | |
| East Java and Madoera..... | June 19-July 3..... | 22 | 1 | Including municipalities in Federal district. |
| Mexico: | | | | |
| Mexico City..... | July 25-31..... | 1 | | |
| Siam: | | | | |
| Bangkok..... | July 11-17..... | 9 | 7 | |

TYPHUS FEVER

| | | | | |
|----------------------|---------------------|----|---|---|
| Chile: | | | | |
| Concepcion..... | June 1-7..... | | 1 | |
| China: | | | | |
| Antung..... | July 26-Aug. 1..... | 3 | | |
| Egypt: | | | | |
| Alexandria..... | July 16-22..... | 1 | | |
| Cairo..... | Feb. 18-25..... | 47 | 7 | |
| Mexico: | | | | |
| Mexico City..... | July 25-31..... | 3 | | Including municipalities in Federal district. |
| Palestine: | | | | |
| Majdal District..... | July 27-Aug. 2..... | 1 | | |
| Yugoslavia..... | July 1-31..... | 2 | 1 | |

YELLOW FEVER

| | | | | |
|------------|----------------|---|--|--|
| Brazil: | | | | |
| Bahia..... | July 4-10..... | 1 | | |

Reports Received from June 26 to September 3, 1926¹**CHOLERA**

| Place | Date | Cases | Deaths | Remarks |
|---------------|-----------------------|-------|--------|--|
| Ceylon..... | | | | Apr. 18-May 29, 1926: Cases, 31; deaths, 29. |
| China: | | | | |
| Shanghai..... | Reported July 20..... | 35 | 8 | |
| Swatow..... | July 11-17..... | | 15 | |

¹ From medical officers of the Public Health Service, American consuls, and other sources.

CHOLERA, PLAGUE, SMALLPOX, TYPHUS FEVER, AND YELLOW FEVER—Continued

Reports Received from June 26 to September 3, 1926—Continued

CHOLERA—Continued

| Place | Date | Cases | Deaths | Remarks |
|-----------------------------|-----------------|-------|--------|---|
| French Settlements in India | | | | Mar. 7-May 15, 1926: Cases, 19; deaths, 18. |
| India | | | | Apr. 26-June 26, 1926: Cases, 18,629; deaths, 11,532. |
| Bombay | May 30-June 5 | 1 | 1 | |
| Calcutta | Apr. 4-May 29 | 478 | 418 | |
| Do. | June 13-26 | 73 | 69 | |
| Do. | June 27-July 10 | 87 | 82 | |
| Madras | May 16-June 5 | 2 | 1 | |
| Rangoon | May 9-June 26 | 67 | 44 | |
| Do. | June 27-July 10 | 16 | 17 | |
| Indo-China: | | | | |
| Saigon | May 2-15 | 52 | 48 | |
| Do. | May 22-June 26 | 42 | 32 | |
| Do. | June 27-July 3 | 19 | 14 | |
| Japan: | | | | |
| Yokohama | Aug. 25 | 1 | | |
| Philippine Islands: | | | | |
| Manila | May 18-24 | 2 | 2 | |
| Do. | June 27-July 17 | 4 | 1 | |
| Provinces— | | | | |
| Albay | Apr. 18-24 | 1 | 1 | |
| Mindoro | Feb. 21-Mar. 6 | 3 | 3 | |
| Romblon | Dec. 14-31 | 42 | 43 | |
| Do. | Jan. 2-23 | 16 | 12 | |
| Siam: | | | | |
| Bangkok | May 2-June 12 | 1,325 | 736 | |
| Do. | June 20-26 | 56 | 26 | |
| Do. | June 27-July 10 | 54 | 22 | |

PLAGUE

| | | | | |
|----------------------|-----------------|----|----|--|
| Algeria: | | | | |
| Algiers | June 21-30 | 1 | | Under date of July 16, 2 cases reported. |
| Azores: | | | | |
| Fayal Island— | | | | |
| Horta | Aug. 2-8 | 1 | 1 | |
| St. Michaels Island | May 9-June 26 | 7 | 2 | |
| British East Africa: | | | | |
| Kisumu | May 16-22 | 1 | 1 | |
| Uganda | Mar. 1-31 | 35 | 34 | |
| Ceylon: | | | | |
| Colombo | May 29-June 5 | 1 | 1 | |
| Chile: | | | | |
| Iquique | June 20-26 | | 1 | |
| China: | | | | |
| Amoy | Apr. 18-June 26 | 40 | 30 | |
| Do. | June 27-July 24 | 21 | | |
| Foochow | June 6-12 | | | Several cases. Not epidemic. |
| Nanking | May 9-July 24 | | | Prevalent. |
| Ecuador: | | | | |
| Guayaquil | May 16-June 30 | 6 | | Rats taken, 36,914; found infected, 31. |
| Do. | July 1-15 | | | Rats taken, 10,020; found infected, 8. |
| Egypt | | | | Jan. 1-July 8, 1925: Cases, 100. |
| City— | | | | |
| Suez | May 21-July 1 | 9 | 5 | |
| Provinces— | | | | |
| Beni-Suef | May 28-June 8 | 8 | 2 | |
| Gharbieh | June 2 | 1 | 1 | |
| France: | | | | |
| Marseille | July 8 | 1 | 1 | Reported July 24. |
| St. Denis | Reported Aug. 2 | 1 | | Vicinity of Paris. |
| Greece: | | | | |
| Athens | Apr. 1-May 31 | 16 | 4 | Including Piræus. |
| Patras | May 27-June 12 | 4 | 1 | |
| Zante | May 17 | 1 | | |
| Hawaii: | | | | |
| Paauhau | July 18-24 | | | Plague-infected rat trapped. |

CHOLERA, PLAGUE, SMALLPOX, TYPHUS FEVER, AND YELLOW FEVER—Continued

Reports Received from June 26 to September 3, 1926—Continued

PLAGUE—Continued

| Place | Date | Cases | Deaths | Remarks |
|------------------------|-----------------|-------|--------|---|
| India | | | | Apr. 25-June 26, 1926: Cases, 53,001; deaths, 41,576. |
| Bombay | May 2-June 26 | 16 | 15 | |
| Karachi | May 23-June 26 | 15 | 13 | |
| Do. | July 11-17 | 1 | 1 | |
| Madras Presidency | Apr. 25-June 26 | 162 | 93 | |
| Do. | July 18-24 | 18 | 12 | |
| Rangoon | May 9-June 26 | 20 | 15 | |
| Do. | June 27-July 10 | 3 | 4 | |
| Indo-China: | | | | |
| Saigon | May 23-June 26 | 8 | 3 | |
| Iraq: | | | | |
| Baghdad | Apr. 18-June 12 | 161 | 108 | |
| Japan: | | | | |
| Yokohama | July 2-30 | 9 | 5 | Total: July 2-Aug. 2, 1926: Cases, 9; deaths, 7. |
| Java: | | | | |
| Batavia | Apr. 24-June 19 | 65 | 65 | |
| Do. | June 26-July 9 | 18 | 17 | |
| Cheribon | Apr. 11-24 | 3 | 3 | |
| East Java and Madoera | June 13-19 | 1 | 1 | |
| Madagascar: | | | | |
| Ambositra Province | May 1-15 | 4 | 4 | Septicemic. |
| Moramanga Province | Apr. 1-15 | 2 | 2 | Do. |
| Tananarive Province | | | | Apr. 1-June 15, 1926: Cases, 120; deaths, 111. |
| Tamatave (Port) | May 16-31 | 1 | 1 | |
| Tananarive Town | Apr. 1-May 15 | 6 | 6 | |
| Other localities | do | 80 | 77 | Bubonic, pneumonic, septicemic. |
| Nigeria | | | | Feb. 1-Apr. 30, 1926: Cases, 115; deaths, 92. |
| Peru | | | | May-June, 1926: Cases, 57; deaths, 16. |
| Departments-- | | | | Present. |
| Ancash | May 1-31 | | | |
| Cajamarca | May 1-June 30 | 10 | 4 | |
| Ica | May 1-31 | 1 | | |
| Libertad | do | 4 | | Pacasmayo, cases, 2; Trujillo district, cases, 2. |
| Lima | May 1-June 30 | 29 | 12 | In Huancabamba district. |
| Piura | June 1-30 | 13 | | Jan. 1-Mar. 31, 1926: Cases, 37. |
| Russia | | | | Nov. 1-30, 1926: Cases, 3; deaths, 2. |
| Senegal | | | | Mar. 1-Apr. 30, 1926: Cases, 15; deaths, 4. |
| Siam: | | | | |
| Bangkok | May 23-June 26 | 2 | 2 | |
| Straits Settlements: | | | | |
| Singapore | May 2-8 | 1 | 1 | |
| Syria: | | | | |
| Beirut | July 1-10 | 1 | | |
| Tunisia: | | | | |
| May 11-June 20 | 150 | | | |
| Kairouan | June 9 | 3 | | 9 cases 30 miles south of Kairouan. |
| Union of South Africa: | | | | |
| Cape Province | May 16-22 | 5 | 3 | |
| Calvinia District | June 13-26 | 12 | 6 | |
| Do. | June 27-July 3 | 1 | | |
| Williston District | June 13-26 | 2 | | |
| Do. | June 27-July 3 | 1 | | |
| Orange Free State-- | | | | |
| Hoopstad District-- | | | | |
| Protestpan | May 9-22 | 3 | 3 | |

SMALLPOX

| | | | |
|----------------|-----------------|-----|----|
| Algeria: | | | |
| Algiers | May 21-June 30 | 14 | |
| Do. | July 1-10 | 1 | |
| Bolivia: | | | |
| La Paz | May 1-June 30 | 14 | 7 |
| Brazil: | | | |
| Bahia | June 20-26 | 1 | |
| Do. | June 27-July 10 | 1 | 7 |
| Manaos | Apr. 1-30 | | 5 |
| Para | May 16-June 26 | 26 | 25 |
| Do. | June 27-July 31 | 14 | 8 |
| Rio de Janeiro | May 2-June 19 | 132 | 91 |
| Santos | Mar. 1-7 | | 1 |

CHOLERA, PLAGUE, SMALLPOX, TYPHUS FEVER, AND YELLOW FEVER—Continued

Reports Received from June 26 to September 3, 1926—Continued

SMALLPOX—Continued

| Place | Date | Cases | Deaths | Remarks |
|-----------------------------|-----------------|-------|--------|---|
| British East Africa: | | | | |
| Tanganyika | May 2-22 | | 12 | |
| Uganda | Mar. 1-31 | 1 | | |
| British South Africa: | | | | |
| Northern Rhodesia | May 18-24 | 17 | 6 | Natives. |
| Do. | June 8-14 | 5 | | |
| Canada | | | | May 30-June 12, 1926: Cases, 46. |
| Alberta | May 30-June 12 | 3 | | |
| Do. | June 27-July 1 | 1 | | |
| Manitoba | May 30-June 26 | 24 | | |
| Do. | June 27-July 24 | 7 | | |
| Winnipeg | June 6-12 | 5 | 1 | |
| Do. | July 4-17 | 6 | | |
| Ontario | | | | May 30-June 25, 1926: Cases, 36. June 27-Aug. 14: Cases, 40. |
| Fort William | July 25-Aug. 7 | 2 | | |
| Kingston | May 23-June 25 | 5 | | |
| Do. | July 11-17 | 2 | 1 | |
| Kitchener | Apr. 26-May 29 | 3 | 1 | |
| North Bay | May 2-22 | 5 | | |
| Do. | July 25-31 | 2 | | |
| Orillia | Apr. 26-May 29 | 7 | | |
| Ottawa | July 18-24 | 1 | | |
| Packenham | do. | 10 | | |
| Toronto | do. | 7 | | |
| Waterloo | do. | 6 | | |
| Saskatchewan | | | | May 30-June 19, 1926: Cases, 16. June 27-Aug. 14: Cases, 37. |
| Regina | July 4-10 | 2 | | Mar. 14-May 29, 1925: Cases, 44; deaths, 3. |
| Ceylon | | | | |
| Chile: | | | | |
| Antofagasta | June 6-12 | 1 | | |
| China: | | | | |
| Amoy | May 1-June 26 | 4 | 8 | |
| Do. | July 4-10 | 1 | | |
| Antung | May 17-June 19 | 5 | | |
| Do. | July 4-18 | 2 | | |
| Canton | May 1-31 | 4 | 2 | |
| Chungking | May 2-July 17 | | | Present. |
| Foochow | May 2-July 10 | | | Do. |
| Hongkong | May 2-June 26 | 19 | 10 | |
| Do. | June 27-July 3 | 1 | 1 | |
| Manchuria | July 6-24 | 13 | | Railway stations. |
| An-shan | May 16-June 12 | 5 | | South Manchurian Railway. |
| Antung | May 19-June 19 | 5 | | |
| Changchun | May 1-June 26 | 6 | | Do. |
| Do. | June 27-July 3 | 1 | | Do. |
| Dairen | Apr. 26-June 20 | 69 | 16 | |
| Do. | June 28-July 18 | 3 | 2 | |
| Fushun | May 16-June 5 | 4 | | Do. |
| Harbin | May 14-June 30 | 21 | | Do. |
| Do. | July 1-21 | 7 | | |
| Kai-yuan | May 16-June 30 | 10 | | Do. |
| Kungehuling | June 13-19 | 1 | | Do. |
| Liao-yang | May 16-June 30 | 4 | | Do. |
| Mukden | do. | 4 | | Do. |
| Penhsihui | May 16-June 19 | 4 | | Do. |
| Ssupingkal | May 16-June 30 | 2 | | Do. |
| Teshihchiao | do. | 2 | | Do. |
| Wa-feng-tien | do. | 3 | | Do. |
| Nanking | May 8-July 24 | | | Present. |
| Shanghai | May 2-June 26 | 10 | 25 | Cases: Foreign. Deaths, popu- |
| Do. | June 27-July 24 | 3 | 3 | lation of international conces- |
| Swatow | May 9-July 10 | | | sion, foreign and native. |
| Tientsia | June 2-26 | | 1 | Sporadic. |
| Wanshien | May 1 | | | Reported by British municip- |
| Chosen | | | | ality. |
| Fusan | May 1-31 | 1 | | Prevalent. |
| Seishun | do. | 2 | 1 | Mar. 1-Apr. 30, 1926: Cases, 368; deaths, 85. |
| Egypt: | | | | |
| Alexandria | May 15-July 1 | 18 | 3 | |
| Cairo | Jan. 29-Feb. 4 | 1 | 1 | |
| Estonia | | | | May 1-June 30, 1926: Cases, 3. |
| France | | | | Mar. 1-Apr. 30, 1926: Cases, 92. |
| St. Etienne | Apr. 18-June 13 | 7 | 2 | |
| French Settlements in India | Mar. 7-May 15 | 205 | 205 | |

CHOLERA, PLAGUE, SMALLPOX, TYPHUS FEVER, AND YELLOW FEVER—Continued

Reports Received from June 26 to September 3, 1926—Continued

SMALLPOX—Continued

| Place | Date | Cases | Deaths | Remarks |
|------------------------------|----------------------|-------|--------|--|
| Gold Coast..... | Mar. 1-Apr. 30.... | 626 | 13 | |
| Great Britain: | | | | |
| England and Wales..... | | | | May 23-July 3, 1926: Cases, 1,068. July 4-31, 1926: Cases, 376. |
| Bradford..... | May 23-29..... | 1 | | |
| Newcastle-on-Tyne..... | June 6-12..... | 1 | | |
| Do..... | July 11-17..... | 1 | | |
| Nottingham..... | May 2-June 5..... | 7 | | |
| Sheffield..... | June 13-19..... | 1 | | |
| Do..... | July 4-10..... | 1 | | |
| Greece: | | | | |
| Saloniki..... | June 1-14..... | | 3 | |
| Guatemala: | | | | |
| Guatemala City..... | June 1-30..... | | 2 | |
| India..... | | | | Apr. 25-June 26, 1926: Cases, 54,831; deaths, 14,771. |
| Bombay..... | May 2-June 26..... | 220 | 134 | |
| Do..... | June 27-July 17..... | 54 | 30 | |
| Calcutta..... | Apr. 4-May 29..... | 171 | 152 | |
| Do..... | June 13-26..... | 24 | 18 | |
| Do..... | June 27-July 10..... | 13 | 12 | |
| Karachi..... | May 16-June 26..... | 44 | 18 | |
| Do..... | June 27-July 24..... | 8 | 4 | |
| Madras..... | May 16-June 26..... | 7 | 4 | |
| Do..... | June 27-July 24..... | 6 | 3 | |
| Rangoon..... | May 9-June 26..... | 10 | 5 | |
| Do..... | July 4-10..... | 1 | | |
| Indo-China: | | | | |
| Saigon..... | do..... | 2 | | |
| Iraq: | | | | |
| Baghdad..... | May 9-June 26..... | 8 | 3 | |
| Do..... | July 4-10..... | 1 | 1 | |
| Basra..... | Apr. 18-June 22..... | 34 | 25 | |
| Italy..... | | | | Mar. 28-June 5, 1926: Cases, 26. Entire consular district, including Island of Sardinia. |
| Rome..... | June 14-20..... | 4 | | Apr. 25-June 26, 1926: Cases, 201. (Reported as alastrim.) |
| Jamaica..... | | | | June 27-July 31, 1926: Cases, 85. (Reported as alastrim.) |
| Do..... | | | | Apr. 11-May 29, 1926: Cases, 564. |
| Japan..... | | | | |
| Kobe..... | May 30-June 5..... | 1 | | |
| Nagoya..... | May 16-22..... | | 1 | |
| Do..... | July 4-10..... | 1 | | |
| Taiwan Island..... | May 11-20..... | 24 | | |
| Do..... | June 1-20..... | 23 | | |
| Tokyo..... | June 26-July 3..... | 2 | | |
| Yokohama..... | May 2-8..... | 2 | | |
| Java: | | | | |
| Batavia..... | May 15-June 25..... | 2 | | Province. |
| East Java and Madoera..... | Apr. 11-June 19..... | 78 | 5 | |
| Malang..... | Apr. 4-10..... | 6 | 1 | Interior. |
| Surabaya..... | May 16-22..... | 14 | 1 | |
| Latvia..... | | | | Apr. 1-30, 1926: Cases, 3. |
| Mexico..... | | | | Feb. 1-Mar. 31, 1926: Deaths, 602. |
| Aguascalientes..... | June 13-26..... | | 5 | |
| Guadalajara..... | June 8-14..... | | 2 | |
| Do..... | June 29-Aug. 16..... | | 5 | |
| Mexico City..... | May 16-June 5..... | 3 | | Including municipalities in Federal District. |
| Saltillo..... | July 18-24..... | | 1 | Present: 100 miles from Chihuahua. |
| San Antonio de Arenales..... | Jan. 1-June 30..... | | | |
| San Luis Potosi..... | June 13-26..... | | 7 | |
| Do..... | July 4-Aug. 14..... | | 9 | |
| Tampico..... | June 1-10..... | | 2 | |
| Torreón..... | May 1-June 30..... | | 17 | |
| Do..... | July 1-31..... | | 5 | |
| Netherlands: | | | | |
| Amsterdam..... | July 18-24..... | | 6 | |
| Nigeria..... | | | | Feb. 1-Apr. 30, 1926: Cases, 404; deaths, 33. |
| Persia: | | | | |
| Teheran..... | Apr. 21-May 21..... | | 7 | |
| Peru: | | | | |
| Arequipa..... | June 1-30..... | | 1 | |
| Poland..... | | | | Mar. 28-May 1, 1926: Cases, 12; deaths, 1. |
| Portugal: | | | | |
| Lisbon..... | Apr. 26-June 19..... | 10 | 3 | |
| Oporto..... | May 23-June 5..... | 4 | | |
| Do..... | July 11-24..... | 2 | | |

CHOLERA, PLAGUE, SMALLPOX, TYPHUS FEVER, AND YELLOW FEVER—Continued

Reports Received from June 26 to September 3, 1926—Continued

SMALLPOX—Continued

| Place | Date | Cases | Deaths | Remarks |
|------------------------|---------------------|-------|--------|---|
| Russia..... | | | | Jan. 1-Mar. 31, 1926: Cases, 2,103. |
| Siam: | | | | |
| Bangkok..... | May 2-June 12..... | 23 | 20 | |
| Do..... | July 4-10..... | 15 | 16 | |
| Straits Settlements: | | | | |
| Singapore..... | Apr. 25-May 1..... | 1 | | |
| Switzerland: | | | | |
| Lucerne Canton..... | June 1-30..... | 1 | | |
| Tunisia..... | | | | Apr. 1-June 30, 1926: Cases, 17. |
| Union of South Africa: | | | | |
| Cape Province..... | June 20-26..... | | | Outbreaks. |
| Idutya district..... | May 23-29..... | | | Do. |
| Orange Free State..... | June 20-July 3..... | | | Do. |
| Natal..... | May 30-June 5..... | | | Do. |
| Transvaal..... | | | | June 6-12, 1926: Outbreaks in Pietersburg and Rustenburg districts. |
| Johannesburg..... | May 9-June 12..... | 5 | | |
| Do..... | July 11-17..... | 1 | | |
| Yugoslavia..... | | | | Apr. 15-30, 1926: Cases, 2; deaths, 1. |
| On vessel..... | | | | Three cases, 1 death, at Aden, Arabia, stated to have been imported by sea. |
| S. S. Karapara..... | | | | At Zanzibar, June 7, 1926. One case of smallpox landed. At Durban, Union of South Africa, June 16, 1926: One suspect case landed. |
| Steamship..... | July 2..... | 1 | | Vessel from Glasgow, Scotland, for Canada. Patient from Glasgow; removed at quarantine on outward voyage. |

TYPHUS FEVER

| | | | | |
|-----------------------------|----------------------|----|---|---|
| Algeria: | | | | |
| Algiers..... | May 21-June 30..... | 7 | 1 | |
| Argentina: | | | | |
| Rosario..... | Feb. 1-23..... | 2 | | |
| Bolivia: | | | | |
| La Paz..... | June 1-30..... | | 1 | |
| Bulgaria..... | | | | Mar. 1-Apr. 30, 1926: Cases, 64; deaths, 12. |
| Chile: | | | | |
| Antofagasta..... | May 23-June 26..... | 4 | | |
| Do..... | June 27-July 3..... | 1 | | |
| Valparaiso..... | Apr. 29-May 5..... | | 1 | |
| China: | | | | |
| Antung..... | June 14-27..... | 7 | 1 | |
| Do..... | June 28-July 18..... | 14 | 1 | |
| Canton..... | May 1-31..... | 1 | | |
| Ichang..... | | | 1 | Reported May 1, 1926. Occurring among troops. |
| Wanshien..... | | | | Present among troops, May 1, 1926. Locality in Chungking consular district. |
| Chosen: | | | | Feb. 1-Apr. 30, 1926: Cases, 640; deaths, 66. |
| Chemulpo..... | May 1-June 30..... | 38 | 2 | |
| Gensan..... | June 1-30..... | 1 | | |
| Seoul..... | do..... | 8 | 3 | |
| Czechoslovakia..... | | | | Jan. 1-May 31, 1926: Cases, 154; deaths, 4. |
| Egypt: | | | | |
| Port Said..... | June 4-24..... | 4 | 1 | |
| Do..... | July 9-15..... | 3 | 1 | |
| Cairo..... | Jan. 29-Feb. 18..... | 8 | 4 | |
| Great Britain: | | | | |
| Scotland— | | | | |
| Glasgow..... | July 30-Aug. 7..... | 9 | | In same family. |
| Ireland (Irish Free State): | | | | |
| Cobh (Queenstown)..... | May 30-June 5..... | 1 | | |
| Do..... | June 27-July 3..... | 1 | 1 | |
| Cork..... | June 5..... | 1 | | |
| Kerry County— | | | | |
| Dingle..... | June 27-July 3..... | 1 | | |

CHOLERA, PLAGUE, SMALLPOX, TYPHUS FEVER, AND YELLOW FEVER—Continued

Reports Received from June 26 to September 3, 1926—Continued

TYPHUS FEVER—Continued

| Place | Date | Cases | Deaths | Remarks |
|-----------------------------|---------------------|-------|--------|---|
| Italy..... | | | | Mar. 28-May 8, 1926: Cases, 3. |
| Japan..... | | | | Mar. 28-May 29, 1926: Cases, 37. |
| Latvia..... | | | | May 1-June 30, 1926: Cases, 19. |
| Lithuania..... | | | | Mar. 1-May 31, 1926: Cases, 172; deaths, 21. |
| Mexico..... | | | | Feb. 1-Mar. 31, 1926: Deaths, 73. |
| Durango..... | July 1-31..... | | 1 | |
| Mexico City..... | May 16-June 5..... | 20 | | Including municipalities in Fed- eral District. |
| Do..... | June 13-19..... | 9 | | Do. |
| San Luis Potosi..... | June 13-26..... | | | Present, city and country. |
| Morocco..... | | | | Mar. 1-May 31, 1926: Cases, 414. |
| Palestine..... | | | | March, 1926: Cases, 6. Exclu- sive of Bedouin tribes and the British military forces. |
| Gaza..... | July 6-12..... | 1 | | |
| Haifa..... | July 13-19..... | 1 | | |
| Jaffa District..... | June 15-28..... | 5 | | |
| Majdal District..... | July 13-19..... | 1 | | |
| Nazareth District..... | do..... | 3 | | |
| Peru:..... | | | | |
| Arequipa..... | Jan. 1-31..... | | 2 | |
| Poland..... | | | | Mar. 29-June 26, 1926: Cases, 1,272; deaths, 85. |
| Rumania..... | | | | Mar. 1-Apr. 30, 1926: Cases, 395; deaths, 49. |
| Russia..... | | | | Jan. 1-Mar. 31, 1926: Cases, 14,814. |
| Tunisia..... | | | | Apr. 1-June 30, 1926: Cases, 110. |
| Tunis..... | June 11-30..... | 3 | | |
| Turkey:..... | | | | |
| Constantinople..... | June 16-22..... | 1 | | |
| Union of South Africa..... | | | | Apr. 1-May 31, 1926: Cases, 153; deaths, 19. |
| Cape Province..... | | | | Apr. 1-May 31, 1926: Cases, 116; deaths, 15. Native. |
| Do..... | May 31-July 3..... | | | Outbreaks. |
| Glengray District..... | June 27-July 3..... | | | Do. |
| Grahamstown..... | do..... | 1 | | Sporadic. |
| Natal..... | | | | Apr. 1-May 31, 1926: Cases, 17. Native. |
| Orange Free State..... | | | | Apr. 1-May 31, 1926: Cases, 15; deaths, 1. |
| Do..... | June 6-12..... | | | Outbreaks. |
| Transvaal..... | | | | Apr. 1-30, 1926: Cases, 3; deaths, 3. Native. |
| Walkkerstroom district..... | June 20-26..... | | | Outbreaks. |
| Wolmaransstad district..... | do..... | | | Do. |
| Yugoslavia..... | | | | Apr. 15-June 30, 1926: Cases, 48; deaths, 7. |
| Zagreb..... | May 15-21..... | 1 | | |

YELLOW FEVER

| Place | Date | Cases | Deaths | Remarks |
|-----------------|-----------------------|-------|--------|---|
| Brazil..... | Reported June 26..... | | | Present in interior of Bahia, Pira- pora, and Minas. |
| Bahia..... | May 9-29..... | 4 | 3 | |
| Do..... | June 6-26..... | 6 | 4 | |
| Gold Coast..... | Apr. 1-10..... | 3 | 1 | |